

# Optimal Approximation of Elliptic Problems II: Wavelet Methods

Stephan Dahlke, Erich Novak, and Winfried Sickel

**1. Introduction.** We study the optimal approximation of the solution of an operator equation

$$(1) \quad \mathcal{A}(u) = f,$$

where  $\mathcal{A}$  is a boundedly invertible linear operator

$$(2) \quad \mathcal{A} : H \rightarrow G$$

from a Hilbert space  $H$  into another Hilbert space  $G$ . We have in mind the more specific situation of an elliptic operator equation, i.e.,

$$(3) \quad \mathcal{A} : H_0^s(\Omega) \rightarrow H^{-s}(\Omega), \quad s > 0,$$

where  $\Omega \subset \mathbf{R}^d$  is a bounded Lipschitz domain. A typical example we shall primary be concerned with is the Poisson equation

$$(4) \quad \begin{aligned} -\Delta u &= f & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega. \end{aligned}$$

Our aim is to answer the following fundamental questions:

- In which sense can we say that an approximation scheme is optimal?
- What happens in the special case of elliptic partial differential equations?
- Do there exist optimal bases and methods, respectively?

**2. Basic Concepts.** We use linear and nonlinear mappings  $S_n$  for the approximation of the solution  $u$  to (1). Let us consider the worst case error

$$e(S_n, F, H) = \sup_{\|f\|_F \leq 1} \|\mathcal{A}^{-1}(f) - S_n(f)\|_H,$$

where  $F$  is a normed (or quasi-normed) space,  $F \subset G$ . For a given basis  $\mathcal{B}$  of  $H$  we consider the class  $\mathcal{N}_n(\mathcal{B})$  of all (linear or nonlinear) mappings of the form

$$S_n(f) = \sum_{k=1}^n c_k h_{i_k},$$

where the  $c_k$  and the  $i_k$  depend in an arbitrary way on  $f$ . Then the nonlinear widths  $e_{n,C}^{\text{non}}(S, F, H)$  are given by

$$e_{n,C}^{\text{non}}(S, F, H) = \inf_{B \in \mathcal{B}_C} \inf_{S_n \in \mathcal{N}_n(\mathcal{B})} e(S_n, F, H).$$

Here  $\mathcal{B}_C$  denotes a set of Riesz bases for  $H$  where  $C$  indicates the stability of the basis. We compare nonlinear approximations with linear approximations. Here we consider the class  $\mathcal{L}_n$  of all continuous linear mappings  $S_n : F \rightarrow H$ ,

$$S_n(f) = \sum_{i=1}^n L_i(f) \cdot \tilde{h}_i$$

with arbitrary  $\tilde{h}_i \in H$ . The worst case error of optimal linear mappings is given by

$$e_n^{\text{lin}}(S, F, H) = \inf_{S_n \in \mathcal{L}_n} e(S_n, F, H).$$

The third class of approximation methods that we study is the class of continuous mappings  $\mathcal{C}_n$ , given by arbitrary continuous mappings  $N_n : F \rightarrow \mathbf{R}^n$  and  $\phi_n : \mathbf{R}^n \rightarrow H$ . Again we define the worst case error of optimal continuous mappings by

$$e_n^{\text{cont}}(S, F, H) = \inf_{S_n \in \mathcal{C}_n} e(S_n, F, H),$$

where  $S_n = \phi_n \circ N_n$ . These numbers, or slightly different numbers, were, e.g., studied by [4, 5, 6]. The three different widths are related as follows:

**Theorem 1.** *Assume that  $S : F \rightarrow H$  with Hilbert spaces  $F$  and  $H$ . Then, under some additional technical conditions*

$$(5) \quad e_n^{\text{lin}}(S, F, H) = e_n^{\text{cont}}(S, F, H) \asymp e_{n, \mathcal{C}}^{\text{non}}(S, F, H).$$

Therefore we conclude that optimal linear mappings have the same order as the best  $n$ -term approximation.

**3. Elliptic Problems.** The next step is to apply this general concept to the special case of elliptic operator equations. It turns out that nonlinear approximation methods do not yield a better rate of convergence compared with linear schemes. The order of convergence only depends on the smoothness of the right-hand side.

**Theorem 2.** *Assume that  $S : H^{-s}(\Omega) \rightarrow H_0^s(\Omega)$  is an isomorphism. Here  $\Omega \subset \mathbf{R}^d$  is a bounded Lipschitz domain. Then for all  $C \geq 1$*

$$(6) \quad e_n^{\text{lin}}(S, H^{-s+t}(\Omega), H^s(\Omega)) \asymp e_{n, \mathcal{C}}^{\text{non}}(S, H^{-s+t}(\Omega), H^s(\Omega)) \asymp n^{-t/d}.$$

However, there is an important difference between regular and nonregular elliptic problems. In the regular case, a Galerkin scheme based on a sequence of uniformly refined spaces is sufficient to obtain the optimal order of convergence, whereas for the nonregular case the optimal linear method requires the precomputation of a suitable basis which is usually a prohibitive task. This leads us to the following problem: Can we find a basis for which best  $n$ -term approximation produces the optimal order of convergence, without any precomputation? We especially focus on a *wavelet basis*  $\Psi = \{\psi_\lambda, \lambda \in \mathcal{J}\}$ . The indices  $\lambda \in \mathcal{J}$  typically encode several types of information, namely the *scale* often denoted  $|\lambda|$ , the spatial location and also the type of the wavelet.  $\Psi$  is assumed to fulfill the following requirements:

- the wavelets are *local* in the sense that

$$\text{diam}(\text{supp}\psi_\lambda) \asymp 2^{-|\lambda|}, \quad \lambda \in \mathcal{J};$$

- the wavelets satisfy the *cancellation property*

$$|\langle v, \psi_\lambda \rangle| \lesssim 2^{-|\lambda|\tilde{m}} \|v\|_{H^{\tilde{m}}(\text{supp}\psi_\lambda)},$$

where  $\tilde{m}$  denotes some suitable parameter;

- the wavelet basis induces characterizations of Besov spaces of the form

$$\|f\|_{B_{\tilde{q}}^s(L_p(\Omega))} \asymp \left( \sum_{|\lambda|=j_0}^{\infty} 2^{j(s+d(\frac{1}{2}-\frac{1}{p}))q} \left( \sum_{\lambda \in \mathcal{J}, |\lambda|=j} |\langle f, \tilde{\psi}_\lambda \rangle|^p \right)^{q/p} \right)^{1/q}, \quad s > d \left( \frac{1}{p} - 1 \right)_+$$

where  $\tilde{\Psi} = \{\tilde{\psi}_\lambda : \lambda \in \mathcal{J}\}$  denotes the *dual basis*,  $\langle \psi_\lambda, \tilde{\psi}_\nu \rangle = \delta_{\lambda,\nu}$ ,  $\lambda, \nu \in \mathcal{J}$ .

It turns out that for the special case of the Poisson equation (4) best  $n$ -term wavelet approximation is still suboptimal, but nevertheless superior when compared with uniform schemes. Moreover, for more specific domains, i.e., for polygon domains, wavelet methods are indeed optimal.

**Theorem 3.** *For the problem (4), best  $n$ -term wavelet approximation produces the worst case error estimate:*

$$(7) \quad e(S_n, H^{t-1}(\Omega), H^1(\Omega)) \leq C n^{-(\frac{t+1}{3}-\varrho)/d} \quad \text{for all } \varrho > 0,$$

provided that  $\frac{1}{2} < t \leq \frac{3d}{2(d-1)} - 1$ .

**Theorem 4.** *For problem (4) in a polygonal domain in  $\mathbf{R}^2$ , best  $n$ -term wavelet approximation is almost optimal in the sense that*

$$(8) \quad e(S_n, H^{t-1}(\Omega), H^1(\Omega)) \leq C n^{-(t-\varrho)/2}, \quad \text{for all } \varrho > 0.$$

The proofs of these results are based on regularity estimates of the exact solution of (4) in specific scales of Besov spaces as developed in [1, 2].

Details of the analysis outlined above can be found in [3].

## References

- [1] S. Dahlke (1999): Besov regularity for elliptic boundary value problems in polygonal domains. *Appl. Math. Lett.* **12(6)**, 31–38.
- [2] S. Dahlke and R. DeVore (1997): Besov regularity for elliptic boundary value problems. *Comm. Partial Differential Equations* **22(1&2)**, 1–16.

- [3] S. Dahlke, E. Novak, and W. Sickel (2004): Optimal approximation of elliptic problems by linear and nonlinear mappings. Bericht Nr. 2004-6, Philipps-Universität Marburg.
- [4] R. A. DeVore, R. Howard, and C. Micchelli (1989): Optimal nonlinear approximation. *Manuscripta Math.* **63**, 469–478.
- [5] R. A. DeVore, G. Kyriazis, D. Leviatan, and V. M. Tikhomirov (1993): Wavelet compression and nonlinear  $n$ -widths. *Adv. Comput. Math.* **1**, 197–214.
- [6] D. Dung and V. Q. Thanh (1996): On nonlinear  $n$ -widths. *Proc. of the AMS* **124**, 2757–276