

Constrained Triangulations, Volumes of Polytopes, and Unit Equations

Michael Kerber¹, Robert Tichy², and Mario Weitzer³

1 Institute of Geometry, Graz University of Technology, Graz, Austria
kerber@tugraz.at

2 Institute of Analysis and Number Theory, Graz University of Technology,
Graz, Austria
tichy@tugraz.at

3 Institute of Analysis and Number Theory, Graz University of Technology,
Graz, Austria
weitzer@math.tugraz.at

Abstract

Given a polytope \mathcal{P} in \mathbb{R}^d and a subset U of its vertices, is there a triangulation of \mathcal{P} using d -simplices that all contain U ? We answer this question by proving an equivalent and easy-to-check combinatorial criterion for the facets of \mathcal{P} . Our proof relates triangulations of \mathcal{P} to triangulations of its “shadow”, a projection to a lower-dimensional space determined by U . In particular, we obtain a formula relating the volume of \mathcal{P} with the volume of its shadow. This leads to an exact formula for the volume of a polytope arising in the theory of unit equations.

1998 ACM Subject Classification F.2.2 [Nonnumerical Algorithms and Problems] Geometrical Problems and Computations, F.2.1 [Numerical Algorithms and Problems] Number-Theoretic Computations

Keywords and phrases constrained triangulations, simplotopes, volumes of polytopes, projections of polytopes, unit equations, S-integers

Digital Object Identifier 10.4230/LIPIcs.SoCG.2017.46

1 Introduction

1.1 Problem statement and results

Let \mathcal{P} be a convex polytope in \mathbb{R}^d , that is, the convex hull of a finite point set V , and let U be a subset of V . We ask for a triangulation of (the interior of) \mathcal{P} with the property that every d -simplex in the triangulation contains all points of U as vertices, calling it a *U-spinal triangulation*. A simple example is the *star triangulation* of \mathcal{P} (Figure 1), where all d -simplices contain a common vertex \mathbf{p} , and U is the singleton set consisting of that point. Another example is the d -hypercube with U being a pair of opposite points (Figure 2). Indeed, the hypercube can be triangulated in a way that all d -simplices contain the space diagonal spanned by U [16, 10].

We are interested in what combinations of \mathcal{P} and U admit spinal triangulations. Our results provide a simple combinatorial answer for this question: Denoting by n the cardinality of U , a U -spinal triangulation of \mathcal{P} exists if and only if each facet of \mathcal{P} contains at least $n - 1$ vertices of U . In that case, we call U a *spine* of \mathcal{P} . More generally, we provide a complete characterization of spinal triangulations: let Φ denote the orthogonal projection of \mathbb{R}^d to the orthogonal complement of the lower-dimensional flat spanned by U . Φ maps U to $\mathbf{0}$ by construction, and \mathcal{P} is mapped to a *shadow* $\hat{\mathcal{P}} := \Phi(\mathcal{P})$. We obtain a U -spinal triangulation



© Michael Kerber, Robert Tichy, and Mario Weitzer;
licensed under Creative Commons License CC-BY


33rd International Symposium on Computational Geometry (SoCG 2017).

Editors: Boris Aronov and Matthew J. Katz; Article No. 46; pp. 46:1–46:15




Leibniz International Proceedings in Informatics


LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



■ **Figure 1** A star triangulation of a hexagon.



■ **Figure 2** A U -spinal triangulation of a cube, where U consists of the two vertices on a space diagonal.



■ **Figure 3** Two examples of the lifting process: (1) Project polytope \mathcal{P} to the orthogonal complement of the flat spanned by U (vertices marked by prominent dots) to obtain shadow $\hat{\mathcal{P}}$. (2) Star-triangulate $\hat{\mathcal{P}}$ with respect to the origin. (3) Lift star triangulation of $\hat{\mathcal{P}}$ to obtain U -spinal triangulation of \mathcal{P} . Note: every facet of \mathcal{P} contains exactly $|U| - 1$ points of U in both examples.

of \mathcal{P} by first star-triangulating $\hat{\mathcal{P}}$ with respect to $\mathbf{0}$ and then lifting each maximal simplex to \mathbb{R}^d by taking the preimage of its vertices under Φ (Figure 3). Vice versa, every spinal triangulation can be obtained in this way.

An important consequence of our characterization is that a spine allows us to relate the volumes of a convex polytope \mathcal{P} and its shadow $\hat{\mathcal{P}}$ (with respect to that spine) by a precise equation. An application of this observation leads to our second result: an exact volume formula of an important polytope arising in number theory which we call the *Everest polytope*. We show that this polytope is the shadow of a higher-dimensional *simplotope*, the product of simplices, whose volume is easy to determine.

1.2 Number theoretic background

In the following we briefly discuss the number-theoretic background of the Everest polytope. G. R. Everest [12, 13] studied various counting problems related to Diophantine equations. In particular, he proved asymptotic results for the number of values taken by a linear form whose variables are restricted to lie inside a given finitely generated subgroup of a number field. This includes norm form- and discriminant form equations, normal integral bases and related objects. Everest’s work contains important contributions to the quantitative theory of S -unit equations and makes use of Baker’s theory of linear forms in logarithms and Schmidt’s subspace theorem from Diophantine approximation; see for instance [30, 14, 2]. Later, other authors [15] applied the methods of Everest to solve combinatorial problems in algebraic number fields. The corresponding counting results involve various important arithmetic constants, one of them being the volume of a certain convex polytope.

In order to introduce Everest's constant, we use basic facts from algebraic number theory. Let K be a number field, $N = N_{K/\mathbb{Q}}$ the field norm and S a finite set of places of K including the archimedean ones. We denote by $O_{K,S} = \{\alpha \in K : |\alpha|_v \leq 1 \text{ for all } v \notin S\}$ the ring of S -integers and its unit group by $U_{K,S}$; the group of S -units. Let c_0, \dots, c_n denote given non-zero algebraic numbers. During the last decades, a lot of work is devoted to the study of the values taken by the expression $c_0x_0 + \dots + c_nx_n$, where the x_n are allowed to run through $U_{K,S}$, see for instance [26, 19]. A specific instance of this kind of general S -unit equations is the following combinatorial problem. As usual, two S -integers α and β are said to be associated (for short $\alpha \sim \beta$) if there exists an S -unit ϵ such that $\alpha = \beta\epsilon$. It is well-known that the group of S -units $U_{K,S}$ is a free abelian group with $s = |S| - 1$ generators, ω_K and $\text{Reg}_{K,S}$ denote as usual the number of roots of unity and the S -regulator of K , respectively (for the basic concepts of algebraic number theory see [25]). Then for given $n \in \mathbb{N}, q > 0$ the counting function $u(n, q)$ is defined as the number of equivalence classes $[\alpha]_{\sim}$ such that

$$N(\alpha) := \prod_{v \in S} |\alpha|_v \leq q, \quad \alpha = \sum_{i=1}^n \varepsilon_i,$$

where $\varepsilon_i \in U_{K,S}$ and no subsum of $\varepsilon_1 + \dots + \varepsilon_n$ vanishes. From the work of Everest [12, 13], the following asymptotic formula can be derived:

$$u(n, q) = \frac{c(n-1, s)}{n!} \left(\frac{\omega_K (\log q)^s}{\text{Reg}_{K,S}} \right)^{n-1} + o((\log q)^{(n-1)s-1+\varepsilon})$$

for arbitrary $\varepsilon > 0$. Here $c(n-1, s)$ is a positive constant, and its exact value has been known only in special cases; see [3] for more details. In general, $c(n, s)$ is given as the volume of a convex polytope in \mathbb{R}^{ns} that we define and study in Section 4. Our results show that

$$c(n, s) = \frac{1}{(s!)^{n+1}} \frac{((n+1)s)!}{(ns)!},$$

which can also be written in terms of a multinomial coefficient as $\binom{(n+1)s}{s, \dots, s} \frac{1}{(ns)!}$.

1.3 Geometric background

Our results fall into the category of *constrained triangulations of convex polytopes*. Triangulations of polytopes are a classic topic in discrete geometry; an infamous question is the quest for triangulating a d -hypercube with a minimal number of simplices [6]. Precise answers are only known up to dimension 7 using computer-assisted proofs [22]. A contemporary overview on results relating to triangulations of polytopes and more general point configurations is provided by de Loera, Rambau, and Santos [8]. It includes a discussion on the triangulation of simplotopes, a geometric object whose study goes back to Hadwiger [20], and has been studied, for instance, in the context of combinatorics [17], game theory [32] and algebraic geometry [18, Ch. 7]. Simplotopes admit a standard triangulation, the so-called *staircase triangulation*, which can easily be described in combinatorial terms. A by-product of our results is that simplotopes can also be triangulated by a family of spinal triangulations.

An n -element subset U of the vertex set of a polytope with the property that each facet contains exactly $n-1$ points of U is called a *special simplex* in the literature. Special simplices have been studied by Athanasiadis [1] to relate the Ehrhart polynomial of integer polytopes with special simplex with the h -vector of the shadow. This results found applications in the study of toric rings and Gorenstein polytopes [21, 5]. Polytopes with special simplices are further studied by de Wolff [9]. His classification yields, among other results, upper bounds

for the number of faces of a polytope with special simplices. Remarkably, special simplices with two vertices, called *spindles*, are also used by Santos for his celebrated counterexample for the Hirsch conjecture [29]. While these works employ similar techniques as our work, for instance, lifting triangulations of the shadow to triangulations of the polytope, the (more elementary) questions of this paper are not addressed in the related work. We also point out that despite the close relation to special simplices, the notion of spines introduced in this paper is slightly more general because a facet is allowed to contain all vertices of U .

Otherwise, constraining triangulations has mostly been considered for low-dimensional problems under an algorithmic angle. For instance, a *constrained Delaunay triangulation* is a triangulation which contains a fixed set of pre-determined simplices; apart from these constraints, it tries to be “as Delaunay as possible”; see Shewchuk’s work [31] for details. While our work is related in spirit, there appears to be no direct connection to this framework, because our constraint does not only ensure the presence of certain simplices in the triangulation, but rather constrains all d -simplices at once.

Computing volumes of high-dimensional convex polytopes is another notoriously hard problem, from a computational perspective [24, Sec. 13][11] as well as for special cases. A famous example is the *Birkhoff polytope* of all doubly-stochastic $n \times n$ -matrices, whose exact volume is known exactly only up to $n = 10$ [27, 7]. Our contribution provides a novel technique to compute volumes of polytopes through lifting into higher dimensions. We point out that lifting increases the dimension, so that the lift of a polytope is not the image of a linear transformation. Therefore, the well-known formula $\text{vol}(A\mathcal{P}) = \sqrt{\det(A^T A)} \text{vol}(\mathcal{P})$ with $A \in \mathbb{R}^{e \times d}$ and $e \geq d$ does not apply to our case.

1.4 Organization

We start by introducing the basic concepts from convex geometry in Section 2. We proceed with our structural result on spinal triangulations, in Section 3. We calculate the vertices of the Everest polytope in Section 4 and define a map from a simplotope to the Everest polytope in Section 5, leading to the volume formula for the Everest polytope. We conclude with some additional remarks in Section 6.

2 Geometric concepts

Let M be an arbitrary subset of \mathbb{R}^d with some integer $d \geq 1$. The *dimension* of M is the dimension of the smallest affine subspace of \mathbb{R}^d containing M . We say that M is *full-dimensional* if its dimension is equal to d . Throughout the entire paper, V will always stand for a finite point set in \mathbb{R}^d that is full-dimensional and in *convex position*, that is $x \notin \text{conv}(V \setminus \{x\})$ for every $x \in V$, where $\text{conv}(\cdot)$ denotes the *convex hull* in \mathbb{R}^d .

2.1 Polytopes and simplicial complexes

We use the following standard definitions (compare, for instance, Ziegler [33]): A *polytope* \mathcal{P} is the convex hull of a finite point set in \mathbb{R}^d in which case we say that the point set *spans* \mathcal{P} . A hyperplane $\mathbb{H} \subseteq \mathbb{R}^d$ is called *supporting* (for \mathcal{P}) if \mathcal{P} is contained in one of the closed half-spaces induced by \mathbb{H} . A *face* of \mathcal{P} is either \mathcal{P} itself, or the intersection of \mathcal{P} with a supporting hyperplane. If a face is neither the full polytope nor empty, we call it *proper*. A face of dimension ℓ is also called ℓ -face of \mathcal{P} , with the convention that the empty set is a (-1) -face. We call the union of all proper faces of \mathcal{P} the *boundary* of \mathcal{P} , and the points of \mathcal{P} not on the boundary the *interior* of \mathcal{P} . 0-faces are called the *vertices* of \mathcal{P} , and we let $V(\mathcal{P})$

denote the set of vertices. With ℓ being the dimension of \mathcal{P} , we call $(\ell - 1)$ -faces *facets*, and $(\ell - 2)$ -faces *ridges* of \mathcal{P} . Any face \mathcal{F} of \mathcal{P} is itself a polytope whose vertex set is $V(\mathcal{P}) \cap \mathcal{F}$.

It is well-known that every point $\mathbf{p} \in \mathcal{P}$ (and only those) can be written as a *convex combination* of vertices of \mathcal{P} , that is $\mathbf{p} = \sum_{\mathbf{v} \in V(\mathcal{P})} \lambda_{\mathbf{v}} \mathbf{v}$ with real values $\lambda_{\mathbf{v}} \geq 0$ for all \mathbf{v} and $\sum_{\mathbf{v} \in V(\mathcal{P})} \lambda_{\mathbf{v}} = 1$. By Carathéodory's theorem, there exists a convex combination with at most $d + 1$ non-zero entries, that is, $\mathbf{p} = \sum_{i=1}^{d+1} \lambda_i \mathbf{v}_i$ with $\mathbf{v}_i \in V(\mathcal{P})$, $\lambda_i \geq 0$ and $\sum \lambda_i = 1$.

An ℓ -simplex σ with $\ell \in \{-1, \dots, d\}$ is a polytope of dimension ℓ that has exactly $\ell + 1$ vertices. Every point in a simplex is determined by a unique convex combination of the vertices. A *simplicial complex* \mathcal{C} in \mathbb{R}^d is a set of simplices in \mathbb{R}^d such that for a simplex σ in \mathcal{C} , all faces of σ are in \mathcal{C} as well, and if σ and τ are in \mathcal{C} , the intersection $\sigma \cap \tau$ is a common face of both (note that the empty set is a face of any polytope). We let $V(\mathcal{C})$ denote the set of all vertices in \mathcal{C} . The *underlying space* $\bigcup \mathcal{C}$ of \mathcal{C} is the union of its simplices. We call a simplex in \mathcal{C} *maximal* if it is not a proper face of another simplex in \mathcal{C} . A simplicial complex equals the set of its maximal simplices together with all their faces and is therefore uniquely determined by its maximal simplices. Also, the underlying space of \mathcal{C} equals the union of its maximal simplices.

In what follows, we let $\mathcal{P} := \text{conv}(V)$ be the polytope spanned by V as fixed above. In particular, $\dim(\mathcal{P}) = d$ because V is full-dimensional and $V(\mathcal{P}) = V$ because V is in convex position.

2.2 Spines

We call $U \subseteq V$ with $|U| = n$ a *spine* of V if each facet of \mathcal{P} contains at least $n - 1$ points of U . Trivially, a one-point subset of V is a spine. If \mathcal{P} is a simplex, any non-empty subset of vertices is a spine. For a hypercube, every pair of opposite vertices forms a spine, but no other spines with two or more elements exist.

We derive a geometric characterization of spines next. The *U -span (in V)* is the set of all d -simplices σ satisfying $U \subseteq V(\sigma) \subseteq V$. Equivalently, it is the set of all d -simplices with vertices in V that have $\text{conv}(U)$ as a common face. Clearly, each simplex of the U -span is contained in \mathcal{P} , and the same is true for the union of all simplices in the U -span.

► **Lemma 1.** *Let $U \subseteq V$ with $|U| = n$. Then, U is a spine of V if and only if the union of all U -span simplices is equal to \mathcal{P} , that is, if every point in \mathcal{P} belongs to at least one simplex in the U -span.*

Proof. We prove both directions of the equivalence separately. For “ \Rightarrow ”, we proceed by induction on n . The statement is true for $n = 0$ by Carathéodory's theorem. Let U be a set with at least one element, $\mathbf{u} \in U$ arbitrary, and $\mathbf{p} \in \mathcal{P} \setminus \{\mathbf{u}\}$. The ray starting in \mathbf{u} through \mathbf{p} leaves the polytope in a point $\bar{\mathbf{p}}$, and this point lies on (at least) one facet \mathcal{F} of \mathcal{P} that does not contain \mathbf{u} . By assumption, \mathcal{F} contains all points in $\bar{U} := U \setminus \{\mathbf{u}\}$. We claim that \bar{U} is a spine of $\bar{V} := V \cap \mathcal{F}$. To see that, note that \mathcal{F} is spanned by \bar{V} and the facets of \mathcal{F} (considered as a polytope in \mathbb{R}^{d-1}) are the ridges of \mathcal{P} contained in \mathcal{F} . Every such ridge \mathcal{R} is the intersection of \mathcal{F} and another facet \mathcal{F}' of \mathcal{P} . By assumption, \mathcal{F}' also contains at least $n - 1$ points of U and it follows at once that \mathcal{R} contains $n - 2$ points of \bar{U} . So, \bar{U} is a spine of \bar{V} , and by induction hypothesis, there exists a $(d - 1)$ -simplex $\bar{\sigma}$ in the \bar{U} -span in \bar{V} that contains $\bar{\mathbf{p}}$. The vertices of $\bar{\sigma}$ together with \mathbf{u} span a simplex σ that contains \mathbf{p} and σ is in the U -span by construction.

The direction “ \Leftarrow ” is clear if $n \in \{0, 1\}$, so we may assume that $n \geq 2$ and proceed by contraposition. If U is not a spine, we have a facet \mathcal{F} of \mathcal{P} such that less than $n - 1$ points

of U lie on \mathcal{F} . Then, every simplex σ in the U -span has at least 2 vertices not on \mathcal{F} , and therefore at most $d-1$ vertices on \mathcal{F} . This implies that $\sigma \cap \mathcal{F}$ is at most $(d-2)$ -dimensional. Therefore, the (finite) union of all U -span simplices cannot cover the $(d-1)$ -dimensional facet \mathcal{F} , which means that the U -span is not equal to \mathcal{P} . ◀

From now on, we use the (equivalent) geometric characterization from the preceding lemma and the combinatorial definition of a spine interchangeably. A useful property is that spines extend to faces in the following sense.

► **Lemma 2.** *Let U be a spine of V , and let \mathcal{F} be an ℓ -face of \mathcal{P} . Then $\bar{U} := U \cap \mathcal{F}$ is a spine of $\bar{V} := V \cap \mathcal{F}$, both considered as point sets in \mathbb{R}^ℓ . In particular, \mathcal{F} contains at least $n - (d - \ell)$ points of U .*

Proof. For every simplex σ in the U -span, let $\bar{\sigma} := \sigma \cap \mathcal{F}$. Clearly, $\bar{\sigma}$ is itself a simplex, spanned by the vertices $V(\bar{\sigma}) = V(\sigma) \cap \mathcal{F}$, and is of dimension at most ℓ . Because U is a spine of V , the union of all $\bar{\sigma}$ covers $\mathcal{F} = \text{conv}(\bar{V})$. Moreover, $V(\bar{\sigma})$ contains \bar{U} . If $\bar{\sigma}$ is not of dimension ℓ , we can find an ℓ -simplex in the \bar{U} -span of \bar{V} that has $\bar{\sigma}$ as a face just by adding suitable vertices from \bar{V} . This implies that the union of the \bar{U} -span covers \mathcal{F} . The “in particular” part follows by downward induction on ℓ . ◀

2.3 Star triangulations

Let $V' \subseteq \mathbb{R}^d$ be a finite point set that is full-dimensional, but not necessarily in convex position. We call a simplicial complex \mathcal{C} a *triangulation* of V' if $V(\mathcal{C}) = V'$ and $\bigcup \mathcal{C} = \text{conv}(V')$. In this case, we also call \mathcal{C} a triangulation of the polytope $\text{conv}(V')$. In a triangulation of V' , every maximal simplex must be of dimension d .

We will consider several types of triangulations in this paper. For the first type, we assume that $V' = V \cup \{\mathbf{0}\}$, where $\mathbf{0} = (0, \dots, 0) \notin V$, V is in convex position (as fixed before), and either $\mathbf{0}$ lies in $\text{conv}(V)$ or V' is in convex position as well. We define a *star triangulation* of V' as a triangulation where all d -simplices contain $\mathbf{0}$ as a vertex. The elementary proof of the following result can be looked up at [23].

► **Lemma 3.** *A star triangulation of V' exists.*

2.4 Pulling triangulations

As usual, let V be a point set in convex position spanning a polytope \mathcal{P} in \mathbb{R}^d , and let $\mathbf{p}_1 \in V$. We can describe a star triangulation with respect to \mathbf{p}_1 also as follows: Triangulate each facet of \mathcal{P} that does not contain \mathbf{p}_1 such that the triangulations agree on their common boundaries. Writing $\Sigma := \{\sigma_1, \dots, \sigma_m\}$ for the maximal simplices triangulating these facets, it is not difficult to see that a (star) triangulation of \mathcal{P} is given by the maximal simplices

$$\mathbf{p}_1 * \Sigma := \{\mathbf{p}_1 * \sigma_1, \dots, \mathbf{p}_1 * \sigma_m\},$$

where $\mathbf{v} * \sigma$ is the simplex spanned by \mathbf{v} and the vertices of σ . Recursively star-triangulating the facets not containing \mathbf{p}_1 in the same way, this construction yields the *pulling triangulation*.

To define the triangulation formally, we fix a total order $\mathbf{p}_1 \prec \mathbf{p}_2 \prec \dots \prec \mathbf{p}_n$ on V . For a single point, we set $\text{Pull}(\{\mathbf{p}\}) := \{\mathbf{p}\}$. For any face \mathcal{F} of \mathcal{P} with positive dimension, let \mathbf{p}_k denote the smallest vertex of \mathcal{F} with respect to \prec . Then

$$\text{Pull}(\mathcal{F}) := \mathbf{p}_k * \bigcup_{\substack{\mathcal{R} \text{ facet of } \mathcal{F} \\ \mathbf{p}_k \notin \mathcal{R}}} \text{Pull}(\mathcal{R}).$$

The following result follows directly by induction on the dimension of the faces. See [4, Sec.5.6], [8, Lemma 4.3.6]:

► **Theorem 4.** *Pull(\mathcal{P}) are the maximal simplices of a triangulation of \mathcal{P} .*

3 Spinal triangulations

Fix a simplicial complex \mathcal{C} with vertex set $V(\mathcal{C})$ in \mathbb{R}^d and a set $U \subseteq V(\mathcal{C})$ of size at most $d + 1$. Let σ denote the simplex spanned by U . We call \mathcal{C} *U-spinal* if every maximal simplex of \mathcal{C} contains σ as a face. If \mathcal{C} is a triangulation of V , we talk about a *U-spinal triangulation* accordingly. *U-spinal triangulations* are closely related to spines: if a *U-spinal triangulation* of V exists, then U is a spine of V , because all maximal simplices of the triangulation lie in the U -span. For the previously discussed spine of a hypercube consisting of two opposite points, also a spinal triangulation exists, consisting of $d!$ d -simplices that all share the diagonal connecting these points. This construction is called *staircase triangulation* [8] or *Freudenthal triangulation* [10].

We show next that a spine always induces a spinal triangulation. Let \mathcal{P} be a polytope spanned by a finite full-dimensional point set $V \subseteq \mathbb{R}^d$ in convex position and let $U = \{\mathbf{u}_1, \dots, \mathbf{u}_n\}$ be a spine of V . We fix a total order \prec on V where $\mathbf{u}_1 \prec \mathbf{u}_2 \prec \dots \prec \mathbf{u}_n$ are the n smallest elements, preceding all points in $V \setminus U$.

► **Lemma 5.** *The pulling triangulation with respect to \prec is a U-spinal triangulation.*

Proof. We prove the statement by induction on n . Let \mathcal{T} denote the pulling triangulation with respect to \prec . For $n = 1$, every maximal simplex of \mathcal{T} contains \mathbf{u}_1 by construction. For $n > 1$, every maximal simplex is a join of \mathbf{u}_1 with a $(d - 1)$ -simplex σ that is contained in a facet \mathcal{F} of \mathcal{P} that does not contain \mathbf{u}_1 . σ , however, is itself a maximal simplex of the pulling triangulation of \mathcal{F} . By the spine property, $U' := \{\mathbf{u}_2, \dots, \mathbf{u}_n\}$ are vertices of \mathcal{F} and form a spine by Lemma 2. By induction, the pulling triangulation of \mathcal{F} is U' -spinal. Therefore, σ contains all vertices of U' , and the join with \mathbf{u}_1 contains all vertices of U . ◀

3.1 Folds and lifts

As before, let \mathcal{P} be a polytope spanned by a finite full-dimensional point set $V \subseteq \mathbb{R}^d$ in convex position, $U \subseteq V$ a spine of V with n elements, and set $e := d - n + 1$. Assume without loss of generality that the origin is among the points in U . Furthermore let \mathbb{A}_U be the subspace of \mathbb{R}^d spanned by U . It is easy to see that the spine points are affinely independent, so that the dimension of \mathbb{A}_U is $n - 1$. Let \mathbb{A}_U^\perp be the orthogonal complement, which is of dimension e . Let $\Phi_U : \mathbb{R}^d \rightarrow \mathbb{A}_U^\perp$ the (orthogonal) projection of \mathbb{R}^d to \mathbb{A}_U^\perp . For notational convenience, we use the short forms $\hat{x} := \Phi_U(x)$ and $\hat{X} := \Phi_U(X)$ for the images of points and sets in \mathbb{R}^d .

Fix a *U-spinal triangulation* \mathcal{T} and let σ be a maximal simplex of \mathcal{T} . Recall that the vertices of σ are the points of U , which all map to $\mathbf{0}$ under Φ_U , and e additional vertices $\mathbf{v}_1, \dots, \mathbf{v}_e$. Hence, $\hat{\sigma}$ is the convex hull of $\{\mathbf{0}, \hat{\mathbf{v}}_1, \dots, \hat{\mathbf{v}}_e\}$. Moreover, since σ has positive (d -dimensional) volume, its projection $\hat{\sigma}$ has positive (e -dimensional) volume as well. It follows that $\hat{\sigma}$ is a e -simplex spanned by $\{\mathbf{0}, \hat{\mathbf{v}}_1, \dots, \hat{\mathbf{v}}_e\}$. Consequently, with $\sigma_1, \dots, \sigma_t$ being the maximal simplices of \mathcal{T} , we call its *fold* the set of simplices consisting of $\hat{\sigma}_1, \dots, \hat{\sigma}_m$ and all their faces. The following statement is a reformulation of [1, Prop.2.3] and [28, Prop.3.12]

► **Lemma 6.** *The fold of a U-spinal triangulation \mathcal{T} is a star triangulation (with respect to the origin).*

Proof. All maximal simplices of the fold contain the origin by construction. Moreover, their union covers $\hat{\mathcal{P}}$ because \mathcal{T} covers \mathcal{P} . Finally, we argue that the fold of two distinct maximal simplices σ_1, σ_2 cannot overlap: Let \mathbb{H} be a hyperplane that separates σ_1 and σ_2 . Since \mathbb{A}_U is contained in the affine span of $\sigma_1 \cap \sigma_2$, \mathbb{H} contains \mathbb{A}_U . Then, $\Phi_U(\mathbb{H})$ is a hyperplane in \mathbb{R}^e which separates the two e -simplices $\hat{\sigma}_1$ and $\hat{\sigma}_2$. \blacktriangleleft

We will now define the converse operation to get from a star triangulation in \mathbb{R}^e to a U -spinal triangulation in \mathbb{R}^d . We first show that pre-images of vertices are well-defined.

► **Lemma 7.** *If $\mathbf{v} \in V \setminus U$ then $\hat{\mathbf{v}} \neq \mathbf{0}$. Furthermore, if $\mathbf{v} \neq \mathbf{w} \in V \setminus U$ then $\hat{\mathbf{v}} \neq \hat{\mathbf{w}}$.*

Proof. Since U is a spine of V , there is a d -simplex σ in the U -span in V which has \mathbf{v} among its vertices. If \mathbf{v} is in the kernel of Φ_U , then \mathbf{v} and the points in U (which span the kernel of Φ_U) are linearly dependent and σ cannot be full-dimensional, which is a contradiction.

For the second part, we assume to the contrary that $\mathbf{v} \neq \mathbf{w}$ but $\hat{\mathbf{v}} = \hat{\mathbf{w}}$. If there is also a d -simplex σ in the U -span which has both \mathbf{v} and \mathbf{w} among its vertices, $\hat{\sigma}$ cannot be full-dimensional, which is a contradiction. Otherwise, if no such d -simplex σ exists, the d -simplices incident to \mathbf{v} triangulate a neighborhood of \mathbf{v} within \mathcal{P} , and the same is true for \mathbf{w} , with the two sets of simplices being disjoint. It follows that their projections under Φ_U have to overlap, contradicting Lemma 6. \blacktriangleleft

For an e -simplex $\hat{\sigma} \subseteq \mathbb{R}^e$ with vertices in \hat{V} and containing $\mathbf{0}$ as vertex, the *lifted* d -simplex $\sigma \subseteq \mathbb{R}^d$ is spanned by the pre-image of $V(\hat{\sigma})$ under $\Phi_U|_V$ (the restriction of Φ_U to V). Note the slight abuse of notation as we chose “ $\hat{\sigma}$ ” as the name of a simplex before even defining the simplex σ , but the naming is justified because $\hat{\sigma}$ indeed is equal to $\Phi_U(\sigma)$ in this case. Given a star triangulation of \hat{V} , its *lift* is given by the set of lifts of its maximal simplices, together with all their faces.

Our goal is to show that the lift of a star triangulation is a U -spinal triangulation of \mathcal{P} . As a first step, we observe that such a lift is a U -spinal simplicial complex.

► **Lemma 8.** *The lift of a star triangulation of \hat{V} is a U -spinal simplicial complex in \mathbb{R}^d whose underlying space is a subset of \mathcal{P} .*

Proof. Fix a star triangulation $\hat{\mathcal{T}}$ and let \mathcal{T} denote its lift. For any simplex in \mathcal{T} , all faces are included by construction. We need to show that for two simplices σ and τ in \mathcal{T} , $\sigma \cap \tau$ is a face of both. We can assume without loss of generality that σ and τ are maximal, hence d -simplices. By construction, σ and τ are the lifts of e -simplices $\hat{\sigma}$ and $\hat{\tau}$ in $\hat{\mathcal{T}}$. Clearly, $\hat{\sigma}$ and $\hat{\tau}$ intersect because they share the vertex $\mathbf{0}$. Moreover, since $\hat{\sigma}$ and $\hat{\tau}$ belong to a triangulation, their intersection is a common face, spanned by a set of vertices $\{\mathbf{0}, \hat{\mathbf{v}}_1, \dots, \hat{\mathbf{v}}_k\}$. Hence, there exists a hyperplane $\hat{\mathbb{H}}$ in \mathbb{R}^e separating $\hat{\sigma}$ and $\hat{\tau}$ such that $\hat{\sigma} \cap \hat{\mathbb{H}} = \text{conv}\{\mathbf{0}, \hat{\mathbf{v}}_1, \dots, \hat{\mathbf{v}}_k\} = \hat{\tau} \cap \hat{\mathbb{H}}$. Let \mathbb{H} denote the preimage of $\hat{\mathbb{H}}$ under Φ_U . Then, \mathbb{H} is a separating hyperplane for σ and τ , and $\sigma \cap \mathbb{H} = \text{conv}\{\mathbf{u}_1, \dots, \mathbf{u}_n, \mathbf{v}_1, \dots, \mathbf{v}_k\} = \tau \cap \mathbb{H}$ as one can readily verify. This shows that the lift is a simplicial complex. Its underlying space lies in \mathcal{P} because every lifted simplex does. It is U -spinal because the lift of every simplex contains U by definition. \blacktriangleleft

3.2 Volumes

The converse of Lemma 6 follows from the fact that all lifts of star triangulations have the same volume, as we will show next.

► **Lemma 9.** *Let σ be a simplex with vertices in V that contains $U = \{\mathbf{u}_1, \dots, \mathbf{u}_n\}$. Then*

$$\binom{d}{n-1} \text{vol}(\sigma) = \text{vol}(U) \text{vol}(\hat{\sigma}),$$

where $\text{vol}(U)$ denotes the volume of the simplex spanned by U .

Proof. For a k -simplex $\tau = \{v_0, \dots, v_k\}$, let $p(\tau)$ denote the parallelotope spanned by $v_1 - v_0, \dots, v_k - v_0$. It is well-known that $\text{vol}(p(\tau)) = k! \text{vol}(\tau)$. Rewriting the claimed volume by expanding the binomial coefficient and noting that $d - (n - 1) = e$ yields

$$\underbrace{d! \text{vol}(\sigma)}_{\text{vol}(p(\sigma))} = \underbrace{(n-1)! \text{vol}(U)}_{\text{vol}(p(U))} \underbrace{e! \text{vol}(\hat{\sigma})}_{\text{vol}(p(\hat{\sigma}))}.$$

To prove the relation between the volumes of parallelotopes, we assume without loss of generality that $\mathbf{u}_1 = \mathbf{0}$. Let \mathbb{A}_U denote the linear subspace spanned by u_2, \dots, u_n , and let $\mathbb{A}_U^{\mathbf{q}}$ denote the parallel affine subspace that contains $\mathbf{q} \in \mathbb{A}_U^\perp$. Then, $\text{vol}(p(\sigma) \cap \mathbb{A}_U^{\mathbf{q}}) = \text{vol}(p(U))$ by definition. By Cavalieri's principle, every parallel cross-section of $p(\sigma)$ has the same volume. More precisely,

$$\text{vol}(p(\sigma) \cap \mathbb{A}_U^{\mathbf{q}}) = \begin{cases} \text{vol}(p(U)) & \text{if } \mathbf{q} \in p(\hat{\sigma}) \\ 0 & \text{otherwise} \end{cases}.$$

Using Fubini's theorem, the volume of $p(\sigma)$ can be expressed as an integral over all cross-sections, which yields

$$\text{vol}(p(\sigma)) = \int_{\mathbf{q} \in \mathbb{A}_U^\perp} \text{vol}(p(\sigma) \cap \mathbb{A}_U^{\mathbf{q}}) d\mathbf{q} = \int_{\mathbf{q} \in p(\hat{\sigma})} \text{vol}(p(U)) d\mathbf{q} = \text{vol}(p(U)) \text{vol}(p(\hat{\sigma})). \quad \blacktriangleleft$$

► **Lemma 10.** *Let U be a spine of V , and \mathcal{T} denote the lift of a star triangulation of \hat{V} . Then, with the notation of Lemma 9,*

$$\binom{d}{n-1} \text{vol}\left(\bigcup \mathcal{T}\right) = \text{vol}(U) \text{vol}(\hat{\mathcal{P}}),$$

where $\bigcup \mathcal{T}$ is the underlying space of \mathcal{T} . In particular, the underlying spaces of the lifts of all star triangulations of \hat{V} have the same volume.

Proof. The statement follows directly from Lemma 9 because the relation holds for any simplex in the star triangulation and its lift. ◀

With that, we can prove our first main theorem.

► **Main Theorem 1 (Lifting theorem).** *Let \mathcal{P} be a polytope spanned by a full-dimensional finite point set $V \subseteq \mathbb{R}^d$ in convex position and $U := \{\mathbf{u}_1, \dots, \mathbf{u}_n\} \subseteq V$. Then,*

U is a spine of V if and only if there exists a U -spinal triangulation of V .

In this case, for $n \geq 1$, the U -spinal triangulations of V are exactly the lifts of the star triangulations of \hat{V} , the orthogonal projection of V to the orthogonal complement of the affine space spanned by U . Furthermore, if $n \geq 2$,

$$\binom{d}{n-1} \text{vol}(\mathcal{P}) = \text{vol}(U) \text{vol}(\hat{\mathcal{P}}).$$

Proof. For the equivalence, the “if”-part follows directly from the geometric characterization of spines (Lemma 1), and the “only if” part follows from Lemma 5.

For the second part, given any U -spinal triangulation \mathcal{T}^* , its fold $\hat{\mathcal{T}}^*$ is a star triangulation by Lemma 6, and by lifting that star triangulation, we obtain \mathcal{T}^* back. Vice versa, starting with any star triangulation $\hat{\mathcal{T}}$, we know that its lift is a simplicial complex contained in \mathcal{P} by Lemma 8. By Lemma 10, the lifts of $\hat{\mathcal{T}}$ and $\hat{\mathcal{T}}^*$ have the same volume, but the lift of the latter is \mathcal{P} . It follows that also the lift of the former is a triangulation of \mathcal{P} , proving the second claim.

The claim about the volumes follows at once by applying Lemma 10 on \mathcal{T}^* and $\hat{\mathcal{T}}^*$. ◀

We remark that not every U -spinal triangulation is a pulling triangulation. This follows from the fact that each pulling triangulation is *regular* (see [8, p.181]), but examples of non-regular spinal triangulations are known (one such example is given in [8, p.306]).

4 The Everest polytope

For $n, s \in \mathbb{N}$, define $\mathcal{E}_{n,s} := \{\mathbf{x} \in \mathbb{R}^{ns} \mid g_{n,s}(\mathbf{x}) \leq 1\}$ where $g_{n,s} : \mathbb{R}^{ns} \rightarrow \mathbb{R}$ with

$$(x_{1,1}, \dots, x_{n,s}) \mapsto \sum_{j=1}^s \max\{0, x_{1,j}, \dots, x_{n,j}\} + \max\left\{0, -\sum_{j=1}^s x_{1,j}, \dots, -\sum_{j=1}^s x_{n,j}\right\}.$$

It is not difficult to verify that $\mathcal{E}_{n,s}$ is bounded and the intersection of finitely many halfspaces of \mathbb{R}^{ns} . We call it the (n, s) -*Everest polytope*. It is well-known [3] that the number-theoretic constant $c(n, s)$ discussed in the introduction is equal to the volume of $\mathcal{E}_{n,s}$.

4.1 Vertex sets

In order to describe the vertices of $\mathcal{E}_{n,s}$ we introduce the following point sets which we also utilize in later parts of the paper. Note that we identify \mathbb{R}^{ns} and $\mathbb{R}^{n \times s}$ which explains the meaning of “row” and “column” in the definition. Let $\mathbf{e}_s(i)$ denote the i -th s -dimensional unit (row) vector with the convention that $\mathbf{e}_s(0) = \mathbf{0}$. We define the following sets in \mathbb{R}^{ns} :

$$V_{n,s} := \left\{ \begin{pmatrix} -\mathbf{e}_s(j_1) \\ \vdots \\ -\mathbf{e}_s(j_n) \end{pmatrix} \mid j_1, \dots, j_n \in \{0, \dots, s\} \right\},$$

$$U_{n,s} := \left\{ \begin{pmatrix} -\mathbf{e}_s(j) \\ \vdots \\ -\mathbf{e}_s(j) \end{pmatrix} \mid j \in \{0, \dots, s\} \right\},$$

$$P_{n,s} := V_{n,s} - (U_{n,s} \setminus \{\mathbf{0}\}) = \{\mathbf{v} - \mathbf{u} \mid \mathbf{v} \in V_{n,s} \wedge \mathbf{u} \in U_{n,s} \setminus \{\mathbf{0}\}\}.$$

It can be readily verified that $V_{n,s}$ is the set of points in $\{-1, 0\}^{ns}$ such that there is at most one -1 per row, $U_{n,s}$ is the set of points in $V_{n,s}$ such that all -1 's (if there are any) are contained in a single column, and $P_{n,s}$ is the set of points in $\{-1, 0, 1\}^{ns}$ such that

- all '1's (if there are any) are in a unique "1-column",
- all entries of the 1-column are either 0 or 1,
- all rows with a 1 contain at most one -1 ,
- all rows without a 1 contain only '0's.

► **Lemma 11.** $P_{n,s} \cap V_{n,s} = \{\mathbf{0}\}$, $U_{n,s} \subseteq V_{n,s}$, $|V_{n,s}| = (s + 1)^n$, $|U_{n,s}| = s + 1$, and $|P_{n,s}| = s(s + 1)^n - s + 1$.

Proof. Follows directly from the definitions and from basic combinatorics. ◀

► **Theorem 12.** *The set of vertices of $\mathcal{E}_{n,s}$ is given by $(P_{n,s} \cup V_{n,s}) \setminus \{\mathbf{0}\} = (V_{n,s} - U_{n,s}) \setminus \{\mathbf{0}\}$.*

We split the proof into several parts which will be treated in the following lemmas. For the rest of this section, let i and j (with a possible subscript) denote elements of $\{1, \dots, n\}$ and $\{1, \dots, s\}$, respectively, let $\mathbf{v} = (v_{1,1}, \dots, v_{n,s})$ be a vertex of $\mathcal{E}_{n,s}$, and set

$$m_j := \max \{0, v_{1,j}, \dots, v_{n,j}\} \text{ for all } j,$$

$$s_i := - \sum_{j=1}^s v_{i,j} \text{ for all } i,$$

$$m := \max \{0, s_1, \dots, s_n\}.$$

Then

$$\begin{aligned} g_{n,s} \begin{pmatrix} v_{1,1} & \cdots & v_{1,s} \\ \vdots & & \vdots \\ v_{n,1} & \cdots & v_{n,s} \end{pmatrix} &= \max \begin{Bmatrix} 0 \\ v_{1,1} \\ \vdots \\ v_{n,1} \end{Bmatrix} + \cdots + \max \begin{Bmatrix} 0 \\ v_{1,s} \\ \vdots \\ v_{n,s} \end{Bmatrix} + \max \begin{Bmatrix} 0 \\ -v_{1,1} - \cdots - v_{1,s} \\ \vdots \\ -v_{n,1} - \cdots - v_{n,s} \end{Bmatrix} \\ &= m_1 + \cdots + m_s + m = 1. \end{aligned}$$

Furthermore it can easily be verified that $v_{i,j} \in [-1, 1]$ for all i and j , $m_j \in [0, 1]$ for all j , $s_i \in [-1, 1]$ for all i , and $m \in [0, 1]$.

In the proofs below, we will repeatedly apply the following argument: if there is an $\varepsilon > 0$ and an $\mathbf{x} \in \mathbb{R}^{ns}$ such that $\mathbf{v} \pm \delta \mathbf{x} \in \mathcal{E}_{n,s}$ for all $\delta \in [0, \varepsilon]$, \mathbf{v} cannot be a vertex of $\mathcal{E}_{n,s}$ (since it is in the interior of an at least 1-dimensional face).

► **Lemma 13.** *If $m_j = 0$ for all j then $\mathbf{v} \in V_{n,s}$.*

Proof. Since all m_j are equal to zero, all $v_{i,j}$ have to be non-positive, so all s_i are non-negative. Also we get that m is equal to 1 which implies that at least one of the s_i is equal to 1. Suppose that $v_{i_0, j_0} \in (-1, 0)$ for some i_0, j_0 . If $s_{i_0} < 1$, then for $\varepsilon := \min \{-v_{i_0, j_0}, 1 - s_{i_0}\} > 0$ and $\delta \in [0, \varepsilon]$ we get that $g_{n,s}(v_{1,1}, \dots, v_{i_0, j_0} \pm \delta, \dots, v_{n,s}) = 1$, so $(v_{1,1}, \dots, v_{i_0, j_0} \pm \delta, \dots, v_{n,s})$ is on the boundary of $\mathcal{E}_{n,s}$ and \mathbf{v} cannot be a vertex of $\mathcal{E}_{n,s}$.

If on the other hand s_{i_0} is equal to 1, then there is a $j_1 \neq j_0$ such that $v_{i_0, j_1} \in (-1, 0)$. But then we get $g_{n,s}(v_{1,1}, \dots, v_{i_0, j_0} \pm \delta, \dots, v_{i_0, j_1} \mp \delta, \dots, v_{n,s}) = 1$ where we set $\varepsilon := \min \{-v_{i_0, j_0}, -v_{i_0, j_1}, v_{i_0, j_0} + 1, v_{i_0, j_1} + 1\} > 0$ and $\delta \in [0, \varepsilon]$, so again \mathbf{v} cannot be a vertex of $\mathcal{E}_{n,s}$.

Thus we get that all $v_{i,j}$ are either -1 or 0 and it is clear that in any given row i_0 only one of the $v_{i_0, j}$ can be -1 (as they sum up to $-s_{i_0} \geq -1$). Therefore $\mathbf{v} \in V_{n,s}$ (see [23] for an extended proof). ◀

► **Lemma 14.** *If $m_{j_0} = 1$ for some j_0 then $\mathbf{v} \in P_{n,s}$.*

Proof. Since m_{j_0} is equal to 1, all other m_j and m have to be equal to zero. Thus all v_{i, j_0} are non-negative and at least one of them is equal to 1. Also, all other $v_{i, j}$ (i.e. if $j \neq j_0$) are non-positive and so are all s_i . Just as in the proof of Lemma 13 we can show that all $v_{i, j}$ are either $-1, 0$, or 1 ; we omit the details. Furthermore it is clear that if v_{i_0, j_0} is equal to 1 for

some i_0 , then there cannot be more than one -1 in the i_0 -th row (as $s_{i_0} \leq 0$). By the same reasoning, if v_{i_0, j_0} is equal to 0, there cannot be any -1 in the i_0 -th row at all. Considering the definition of $P_{n,s}$ we thus see that $\mathbf{v} \in P_{n,s}$. ◀

► **Lemma 15.** *If $m_j \neq 1$ for all j then $m_j = 0$ for all j .*

Proof. The proof works similar to that of Lemma 13; see [23]. ◀

Proof of Theorem 12. Lemma 15 implies that if \mathbf{v} is a vertex of $\mathcal{E}_{n,s}$, then we are in the situation of either Lemma 13 or Lemma 14, hence $\mathbf{v} \in P_{n,s} \cup V_{n,s}$. Furthermore it is clear that $\mathbf{0}$ is not a vertex of $\mathcal{E}_{n,s}$. Also, it follows from the definition of $P_{n,s}$ that

$$E_{n,s} := (P_{n,s} \cup V_{n,s}) \setminus \{\mathbf{0}\} = ((V_{n,s} - (U_{n,s} \setminus \{\mathbf{0}\})) \cup V_{n,s}) \setminus \{\mathbf{0}\} = (V_{n,s} - U_{n,s}) \setminus \{\mathbf{0}\}.$$

We are left to show that $E_{n,s} \subseteq V(\mathcal{E}_{n,s})$. First we observe that $g_{n,s}(\mathbf{v}) = 1$ for all $\mathbf{v} \in E_{n,s}$. We consider the case that $n, s \geq 2$ and assume that there is a $\mathbf{v} \in E_{n,s}$ that is not a vertex of $\mathcal{E}_{n,s}$. Since \mathbf{v} is on the boundary of $\mathcal{E}_{n,s}$ but not a vertex of $\mathcal{E}_{n,s}$, it is contained in the interior of an at least 1-dimensional face \mathcal{F} of $\mathcal{E}_{n,s}$. Let \mathbf{w} be any vertex of \mathcal{F} . Then $\mathbf{w} \in E_{n,s}$ and $\mathbf{v} \neq \mathbf{w}$.

Now let $\mathbf{a} \in E_{2,2}$, $\mathbf{a} \neq \mathbf{b} \in E_{2,2} \cup \{\mathbf{0}\}$, and consider the convex combination $\alpha\mathbf{a} + (1-\alpha)\mathbf{b}$, $\alpha \in \mathbb{R}$. By plugging in all possible values of \mathbf{a} and \mathbf{b} one can verify that if $\alpha > 1$ then $g_{2,2}(\alpha\mathbf{a} + (1-\alpha)\mathbf{b}) > 1$.

Let $\mathbf{v}', \mathbf{w}' \in \mathbb{R}^{2 \times s}$ be submatrices consisting of 2 rows of \mathbf{v} and \mathbf{w} respectively, such that $\mathbf{v}' \neq \mathbf{0}$ and $\mathbf{v}' \neq \mathbf{w}'$. By definition of $E_{n,s}$, \mathbf{v}' and \mathbf{w}' thus respectively contain submatrices of the form \mathbf{a} and \mathbf{b} from above while the remaining entries are padded with zeros. It follows from the definition of $g_{n,s}$ that $g_{n,s}(\alpha\mathbf{v}' + (1-\alpha)\mathbf{w}') > 1$ if $\alpha > 1$, which contradicts the fact that \mathbf{v} is in the interior of \mathcal{F} . Hence, $E_{n,s} \subseteq V(\mathcal{E}_{n,s})$ if $n, s \geq 2$. If $n = 1$ and $s \geq 2$ one can proceed analogously by considering $\mathbf{a} \in E_{1,2}$, $\mathbf{a} \neq \mathbf{b} \in E_{1,2} \cup \{\mathbf{0}\}$; same goes for $s = 1$ and $\mathbf{a} \in E_{1,1}$, $\mathbf{a} \neq \mathbf{b} \in E_{1,1} \cup \{\mathbf{0}\}$. ◀

► **Corollary 16.** *The number of vertices of $\mathcal{E}_{n,s}$ is given by $(s+1)^{n+1} - s - 1$.*

Proof. Follows directly from Theorem 12 and Lemma 11. ◀

5 Projections of simplotopes

We will establish a relation between the Everest polytope $\mathcal{E}_{n,s}$ and a special polytope known as simplotope. This relation will allow the comparison of the volumes of the two polytopes even though they are of different dimension.

5.1 Simplotopes

For $s \in \mathbb{N}$, the s -simplex Δ_s is spanned by the points $(\mathbf{0}, -\mathbf{e}_s(1), \dots, -\mathbf{e}_s(s))$ in \mathbb{R}^s , with $\mathbf{e}_s(i)$ the i -th standard vector in \mathbb{R}^s , as before. A *simplotope* is a Cartesian product of the form $\Delta_{s_1} \times \dots \times \Delta_{s_n}$ with positive integers s_1, \dots, s_n . Note that in the literature, simplotopes are usually defined in a combinatorially equivalent way using the standard s -simplex spanned by $(s+1)$ -unit vectors in \mathbb{R}^{s+1} . We restrict to the case that all s_i are equal, and we call

$$\mathcal{S}_{n,s} = \underbrace{\Delta_s \times \dots \times \Delta_s}_{n \text{ times}}$$

the (n, s) -simplotope for $n, s \in \mathbb{N}$.

For instance, d -hypercubes are d -fold products of line segments, and therefore (n, s) -simplotopes with $n = d$ and $s = 1$. It is instructive to visualize a point in $\mathcal{S}_{n,s}$ as an $n \times s$ -matrix with real entries in $[0, 1]$, where the sums of the entries in each row do not exceed 1. One can readily verify that the set of vertices of the (n, s) -simplotope is equal to $V_{n,s}$, as given in the beginning of Section 4. Moreover, it is straight-forward to verify that each facet of $\mathcal{S}_{n,s}$ is given by

$$\underbrace{\Delta_s \times \dots \times \Delta_s}_i \times \mathcal{F} \times \underbrace{\Delta_s \times \dots \times \Delta_s}_{n-i-1 \text{ times}}$$

where \mathcal{F} is a facet of Δ_s and $0 \leq i \leq n - 1$. It follows at once:

► **Theorem 17.** $U_{n,s}$ is a spine of $V_{n,s} = V(\mathcal{S}_{n,s})$.

5.2 A linear transformation

We call the matrix

$$\Pi_{n,s} := \begin{pmatrix} & -I_s \\ I_{ns} & \vdots \\ & -I_s \end{pmatrix} \in \mathbb{R}^{(ns) \times ((n+1)s)},$$

where I_d is the identity matrix of dimension d , the (n, s) -SE-transformation (“SE” stands for “Simplotope \leftrightarrow Everest polytope”). We show that the name is justified, as it maps the $(n + 1, s)$ -simplotope onto the (n, s) -Everest polytope. See [23] for a proof.

► **Theorem 18.** $\Pi_{n,s}(V(\mathcal{S}_{n+1,s}) \setminus U_{n+1,s}) = V(\mathcal{E}_{n,s})$, $U_{n+1,s} \setminus \{\mathbf{0}\}$ is a basis of $\ker(\Pi_{n,s})$ (in particular, $\Pi_{n,s}U_{n+1,s} = \{\mathbf{0}\}$), and $\mathbf{0}$ is contained in the interior of $\mathcal{E}_{n,s}$. In particular, $\Pi_{n,s}\mathcal{S}_{n+1,s} = \mathcal{E}_{n,s}$.

We are now ready to prove a formula for the Everest polytope.

► **Main Theorem 2.** The volume of the (n, s) -Everest polytope is given by

$$\text{vol}(\mathcal{E}_{n,s}) = \frac{((n + 1)s)!}{(ns)!(s!)^{n+1}}.$$

Proof. We want to apply the Lifting Theorem (Main Theorem 1) with $\mathcal{P} \leftarrow \mathcal{S}_{n+1,s}$, $\hat{\mathcal{P}} \leftarrow \mathcal{E}_{n,s}$, $d \leftarrow (n + 1)s$ and $e \leftarrow ns$. However, a minor modification is needed, because the SE-transformation $\Pi_{n,s}$ is not a projection matrix. So, let $\tilde{\Pi}_{n,s} := \begin{pmatrix} \Pi_{n,s} \\ 0 \ I_s \end{pmatrix} \in \mathbb{R}^{((n+1)s) \times ((n+1)s)}$,

$\Pi := (I_{ns} \ 0) \in \mathbb{R}^{(ns) \times ((n+1)s)}$, and let $\tilde{\mathcal{S}}_{n+1,s} := \tilde{\Pi}_{n,s}\mathcal{S}_{n+1,s}$ denote the transformed simplotope. Clearly, $\text{vol}(\tilde{\mathcal{S}}_{n+1,s}) = \text{vol}(\mathcal{S}_{n+1,s})$, $\Pi\tilde{\mathcal{S}}_{n+1,s} = \mathcal{E}_{n,s}$, and the transformed spine points $\tilde{U}_{n+1,s} := \tilde{\Pi}_{n,s}U_{n+1,s}$ span the kernel of Π . Using Main Theorem 1 on $\tilde{\mathcal{S}}_{n+1,s}$ and $\mathcal{E}_{n,s}$, we obtain

$$\binom{(n + 1)s}{ns} \text{vol}(\tilde{\mathcal{S}}_{n+1,s}) = \text{vol}(\tilde{U}_{n+1,s}) \text{vol}(\mathcal{E}_{n,s}).$$

Furthermore, $\text{vol}(\tilde{U}_{n+1,s}) = \frac{1}{s!}$ since $\tilde{U}_{n+1,s} = \{(\mathbf{0}, \dots, \mathbf{0}, -\mathbf{e}_s(j)) \mid j \in \{0, \dots, s\}\}$. Moreover, $\mathcal{S}_{n+1,s}$ is the $(n + 1)$ -fold product of simplices Δ_s spanned by $\mathbf{0}$ and s unit vectors. Hence, the volume of Δ_s is $1/s!$, and by Fubini’s theorem,

$$\text{vol}(\tilde{\mathcal{S}}_{n+1,s}) = \text{vol}(\mathcal{S}_{n+1,s}) = (\text{vol}(\Delta_s))^{n+1} = \frac{1}{(s!)^{n+1}}.$$

Plugging in the formulas for $\text{vol}(\tilde{U}_{n+1,s})$ and $\text{vol}(\tilde{\mathcal{S}}_{n+1,s})$ into the formula given by the lifting theorem the claim follows by rearranging terms. ◀

6 Conclusions and further remarks

Main Theorem 1 combines several new results. It answers the question of the existence of a triangulation of a polytope under the constraint that a given subset of the vertices of the polytope must be contained in every maximal simplex of the triangulation. Furthermore, it characterizes all such triangulations and provides a method to compute one (or all) efficiently from the lift of a star triangulation. Finally, it generalizes the well-known relation $\text{vol}(AM) = |\det(A)| \text{vol}(M)$ where M is a measurable subset of \mathbb{R}^d and $A \in \mathbb{R}^{d \times d}$ to certain cases where A is not a square matrix. In particular, it allows us to express the volume of an object in \mathbb{R}^d in terms of the volume its “shadow” in \mathbb{R}^e , and vice versa.

The shadow that a cube in \mathbb{R}^3 casts if the light shines parallel to any of its space diagonals is a regular hexagon. Assuming a cube with side length ℓ , the theorem implies that the volume of the cube and the volume (area) of its shadow (the hexagon) differ by a factor of $\sqrt{3}/\ell$ which provides an alternative method to compute the volume of a hexagon from the volume of a cube. By lifting the “complicated” hexagon to a higher dimensional space it gains more symmetries and becomes the comparatively simple cube. In the same fashion, the complicated Everest polytope is the shadow of the simpler simplotope which allowed the computation of its volume in Main Theorem 2.

Starting with a polytope and a spine, it is easy to determine the volume of the “shadow” with respect to the spine using our theorem. On the other hand, there is no easy way to tell if a given shape is the shadow of some higher dimensional object and in the case of the Everest polytope, this is the interesting direction. We pose the question of whether other polytopes (e.g., the Birkhoff polytope) can be expressed as shadows of other polytopes. For that purpose, it might be worthwhile to find general methods or at least good heuristics to determine if a complicated shape can be recognized as the shadow of some simpler object.

Acknowledgements. We thank Raman Sanyal, Volkmar Welker, and the anonymous referees for helpful discussions and recommendations that have led to significant simplifications of our exposition.

References

- 1 C. Athanasiadis. Ehrhart polynomials, simplicial polytopes, magic squares and a conjecture of Stanley. *J. Reine Angew. Math.*, 583:163–174, 2005.
- 2 A. Baker and G. Wüstholz. Logarithmic forms and group varieties. *J. Reine Angew. Math.*, 442:19–62, 1993.
- 3 F. Barroero, C. Frei, and R. Tichy. Additive unit representations in rings over global fields – a survey. *Publ. Math. Debrecen*, 79(3–4):291–307, 2011.
- 4 M. Beck and R. Sanyal. *Combinatorial Reciprocity Theorems*. American Mathematical Society, 2016. In preparation, available at <http://math.sfsu.edu/beck/crt.html>.
- 5 W. Bruns and T. Römer. h-vectors of Gorenstein polytopes. *J. Combin. Theory Ser. A*, 114:65–76, 2007.
- 6 H. Croft, K. Falconer, and R. Guy. *Unsolved Problems in Geometry*. Springer, 1991.
- 7 J. de Loera, F. Liu, and R. Yoshida. A generating function for all semi-magic squares and the volume of the Birkhoff polytope. *J. Algebraic Combin.*, pages 113–139, 2009.
- 8 J. de Loera, J. Rambau, and F. Santos. *Triangulations*. Springer, 2010.
- 9 T. de Wolff. Polytopes with special simplices. arXiv:1009.6158.
- 10 H. Edelsbrunner and M. Kerber. Dual complexes of cubical subdivisions of \mathbb{R}^n . *Discrete Comput. Geom.*, 47(2):393–414, 2012.

- 11 I. Emiris and V. Fisikopoulos. Efficient random-walk methods for approximating polytope volume. In *Proc. of the 13th Annual Symp. on Comp. Geom.*, SOCG'14, pages 318:318–318:327, 2014.
- 12 G. R. Everest. A “Hardy-Littlewood” approach to the S -unit equation. *Compos. Math.*, 70(2):101–118, 1989.
- 13 G. R. Everest. Counting the values taken by sums of S -units. *J. Number Theory*, 35(3):269–286, 1990.
- 14 J. Evertse and H. P. Schlickewei. A quantitative version of the absolute subspace theorem. *J. Reine Angew. Math.*, 548:21–127, 2002.
- 15 C. Frei, R. Tichy, and V. Ziegler. On sums of S -integers of bounded norm. *Monatsh. Math.*, 175(2):241–247, 2014.
- 16 H. Freudenthal. Simplicialzerlegung beschränkter Flachheit. *Ann. of Math.*, pages 580–582, 1942.
- 17 R. Freund. Combinatorial theorems on the simplotope that generalize results on the simplex and cube. *Math. Oper. Res.*, 11(1):169–179, 1986.
- 18 I. Gelfand, M. Kapranov, and A. Zelevinsky. *Discriminants, Resultants and Multidimensional Determinants*. Birkhäuser, 2008.
- 19 K. Györy and K. Yu. Bounds for the solutions of S -unit equations and decomposable form equations. *Acta Arith.*, 123(1):9–41, 2006.
- 20 H. Hadwiger. *Vorlesungen über Inhalt, Oberfläche und Isoperimetrie*. Springer, 1957.
- 21 T. Hibi and H. Ohsugi. Special simplices and Gorenstein toric rings. *J. Combin. Theory Ser. A*, 113:718–725, 2006.
- 22 R. Hughes and M. Anderson. Simplexity of the cube. *Discrete Math.*, 158:99–150, 1996.
- 23 M. Kerber, R. Tichy, and M. Weitzer. Constrained triangulations, volumes of polytopes, and unit equations. *arXiv*, 1609.05017, 2016.
- 24 J. Matoušek. *Lectures in Discrete Geometry*. Springer, 2002.
- 25 J. Neukirch. *Algebraic number theory*, volume 322 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, 1999.
- 26 K. Nishioka. Algebraic independence by Mahler’s method and S -unit equations. *Compositio Math.*, 92(1):87–110, 1994.
- 27 I. Pak. Four questions on Birkhoff polytope. *Ann. Comb.*, 4:83–90, 2000.
- 28 V. Reiner and V. Welker. On the Charney-Davis and Neggers-Stanley conjectures. *J. Combin. Theory Ser. A*, 109(2):247–280, 2005.
- 29 F. Santos. A counterexample to the Hirsch conjecture. *Ann. of Math.*, 176:383–412, 2012.
- 30 H. P. Schlickewei. S -unit equations over number fields. *Invent. Math.*, 102(1):95–108, 1990.
- 31 J. Shewchuk. General-dimensional constrained Delaunay and constrained regular triangulations, I: Combinatorial properties. *Discrete Comput. Geom.*, 39:580–637, 2008.
- 32 G. van der Laan and A. Talman. On the computation of fixed points in the product space of unit simplices and an application to noncooperative n person games. *Math. Oper. Res.*, 7(1):1–13, 1982.
- 33 G. Ziegler. *Lectures on Polytopes*. Springer, 2007.