

How Does Object Fatness Impact the Complexity of Packing in d Dimensions?

Sándor Kisfaludi-Bak

Max Planck Institut für Informatik, Saarbrücken, Germany
sandor.kisfaludi-bak@mpi-inf.mpg.de

Dániel Marx

Max Planck Institut für Informatik, Saarbrücken, Germany
dmarx@cs.bme.hu

Tom C. van der Zanden

Department of Data Analytics and Digitalisation, Maastricht University, The Netherlands
T.vanderZanden@maastrichtuniversity.nl

Abstract

Packing is a classical problem where one is given a set of subsets of Euclidean space called objects, and the goal is to find a maximum size subset of objects that are pairwise non-intersecting. The problem is also known as the Independent Set problem on the intersection graph defined by the objects. Although the problem is NP-complete, there are several subexponential algorithms in the literature. One of the key assumptions of such algorithms has been that the objects are fat, with a few exceptions in two dimensions; for example, the packing problem of a set of polygons in the plane surprisingly admits a subexponential algorithm. In this paper we give tight running time bounds for packing similarly-sized non-fat objects in higher dimensions.

We propose an alternative and very weak measure of fatness called the stabbing number, and show that the packing problem in Euclidean space of constant dimension $d \geq 3$ for a family of similarly sized objects with stabbing number α can be solved in $2^{O(n^{1-1/d}\alpha)}$ time. We prove that even in the case of axis-parallel boxes of fixed shape, there is no $2^{o(n^{1-1/d}\alpha)}$ algorithm under ETH. This result smoothly bridges the whole range of having constant-fat objects on one extreme ($\alpha = 1$) and a subexponential algorithm of the usual running time, and having very “skinny” objects on the other extreme ($\alpha = n^{1/d}$), where we cannot hope to improve upon the brute force running time of $2^{O(n)}$, and thereby characterizes the impact of fatness on the complexity of packing in case of similarly sized objects. We also study the same problem when parameterized by the solution size k , and give a $n^{O(k^{1-1/d}\alpha)}$ algorithm, with an almost matching lower bound: there is no algorithm with running time of the form $f(k)n^{o(k^{1-1/d}\alpha/\log k)}$ under ETH. One of our main tools in these reductions is a new wiring theorem that may be of independent interest.

2012 ACM Subject Classification Theory of computation \rightarrow Computational geometry

Keywords and phrases Geometric intersection graph, Independent Set, Object fatness

Digital Object Identifier 10.4230/LIPIcs.ISAAC.2019.36

Related Version A full version of the paper is available at <http://arxiv.org/abs/1909.12044>.

Funding *Sándor Kisfaludi-Bak*: Supported by NWO Gravitation Grant Networks (No. 024.002.003).
Dániel Marx: Supported by ERC Consolidator Grant SYSTEMATICGRAPH (No. 725978).

1 Introduction

Many well-known NP-hard problems (e.g. INDEPENDENT SET, HAMILTON CYCLE, DOMINATING SET) can be solved in time $2^{O(\sqrt{n})}$ when restricted to planar graphs, while only $2^{O(n)}$ algorithms are known for general graphs [11–16, 18, 24, 28, 30]. This beneficial effect of planarity is known as the “square root phenomenon,” and can be exploited also in the context of 2-dimensional geometric problems where the problem is defined on various intersection



© Sándor Kisfaludi-Bak, Dániel Marx, and Tom C. van der Zanden;
licensed under Creative Commons License CC-BY

30th International Symposium on Algorithms and Computation (ISAAC 2019).

Editors: Pinyan Lu and Guochuan Zhang; Article No. 36; pp. 36:1–36:18

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

graphs in \mathbb{R}^2 [3, 4, 17, 25]. In particular, consider the geometric packing problem where, given a set of polygons in \mathbb{R}^2 , the task is to find a subset of k pairwise disjoint polygons. This problem can be solved in time $n^{O(\sqrt{k})}$ [25], which – when expressed only as a function of the input – gives an $n^{O(\sqrt{n})} = 2^{O(\sqrt{n} \log n)}$ algorithm for finding a maximum size disjoint subset.

Can these 2-dimensional subexponential algorithms be generalized to higher dimensions? It seems that the natural generalization is to aim for $2^{O(n^{1-1/d})}$, or in case of parameterized problems, either $2^{O(k^{1-1/d})} \cdot n^{O(1)}$ or $n^{O(k^{1-1/d})}$ time algorithms in d -dimensions: the literature contains upper and lower bounds of this form (although sometimes with extra logarithmic factors in the exponent) [9, 26, 29]. However, all of these algorithms have various restrictions on the object family on which the intersection graph is based: there is no known analogue of the $n^{O(\sqrt{k})}$ time algorithm of Marx and Pilipczuk [25] in higher dimensions with the same generality of objects. There is a good reason for this: it is easy to see that any n -vertex graph can be expressed as the intersection graph of 3-dimensional simple polyhedra. Thus a subexponential algorithm for 3-dimensional objects without any severe restriction would give a subexponential algorithm for INDEPENDENT SET on general graphs, violating standard complexity-theoretic assumptions.

What could be reasonable restrictions on the objects that allow running times of the form, e.g., $2^{O(n^{1-1/d})}$? One of the most common restrictions is to study a set $F \subseteq 2^{\mathbb{R}^d}$ of *fat objects*, where for each object $o \in F$ the ratio $\text{radius}(B_{\text{in}}(o)) / \text{radius}(B_{\text{out}}(o))$ is at least some fixed positive constant. (We denote by $\text{radius}(B_{\text{in}})$ and $\text{radius}(B_{\text{out}})$ the radius of the inscribed and circumscribed ball respectively.) Another common restriction is to have *similarly sized* objects, that is, a family F where the ratio of the largest and smallest object diameter is at most some absolute constant. Many results concern only *unit disk graphs*, where F consists of unit disks in the plane: unit disks are both fat and similarly sized. The focus of our paper is to explore the role of fatness in the context of packing problems and to understand when and to what extent fatness decreases the complexity of the problem. We observe that fatness is a crucial requirement for subexponential algorithms in higher dimensions, and this prompts us to explore in a quantitative way how fatness influences the running time. For this purpose, we introduce a parameter α describing the fatness of the objects and give upper and lower bounds taking into account this parameter as well.

More precisely, we introduce the notion of the *stabbing number*, which can be regarded as an alternative measure of fatness. This slightly extends a similar definition by Chan [6]. We say that an object o is stabbed by a point p if $p \in o$. A family of objects $F \subseteq 2^{\mathbb{R}^d}$ is α -stabbed if for any $r \in \mathbb{R}$, the subset of F -objects o of diameter $\text{diam}(o) \in [r/2, r)$ contained in any ball of radius r can be stabbed by α^d points. The stabbing number of F is defined as $\inf_{\alpha \in [1, \infty)} \{F \text{ is } \alpha\text{-stabbed}\}$. Note that a set of n objects in d -dimensions has stabbing number at most $n^{1/d}$. The stabbing number is closely related to the inverse of a common measure of fatness. This relationship is explored in Section 2.

By adapting a separator theorem from [9], we can give an algorithm where the running time smoothly goes from $2^{O(n^{1-1/d})}$ to $2^{O(n)}$ as the stabbing number goes from $O(1)$ to the maximum possible $n^{1/d}$.

► **Theorem 1.** *Let $\alpha \in [1, \infty)$ and $2 \leq d \in \mathbb{N}$ be fixed constants. There is an algorithm that solves INDEPENDENT SET for intersection graphs of similarly sized α -stabbed objects in \mathbb{R}^d running in time $2^{O(n^{1-1/d} \alpha)}$.*

As mentioned, the stabbing number is at most $n^{1/d}$, and this algorithm runs in subexponential time whenever the stabbing number is better than this trivial upper bound, that is, whenever $\alpha = o(n^{1/d})$ holds.

In order to have definite answers to the best running times achievable, we also need a lower bound framework. A popular starting point in the past decades is the Exponential Time Hypothesis (ETH) [21], which posits that there exists a constant $\gamma > 0$ such that there is no $2^{\gamma n}$ algorithm for the 3-SAT problem. Classical NP-hardness reductions automatically yield quantitative lower bounds on the running time under ETH. If enough care is taken to ensure that the constructed instance is sufficiently small, then one can find lower bounds that match the best known algorithms [8]. For the INDEPENDENT SET problem, a lower bound of $2^{\Omega(n)}$ is a consequence of classical reductions under ETH.

A standard way to explore the impact of a parameter such as fatness is to give an algorithm where the parameter appears in the running time, together with a matching lower bound. However, the notion of “matching lower bound” needs to be defined precisely if we are expressing the running time as a function of two parameters, the size n of the instance and the stabbing number α of the objects.

A recent example of such an algorithm and lower bound involving two parameters is the paper by Biró *et al.* [5], where it is shown that the coloring problem of unit disk graphs with $\ell = n^\lambda$ colors can be solved in $2^{O(\sqrt{n\ell} \log n)}$ time, where $\lambda \in [0, 1]$ is a fixed constant, and they also exclude algorithms of running time $2^{o(\sqrt{n\ell})}$ under ETH. This is interesting since this smoothly bridges the gap between a standard “square root phenomenon” algorithm ($\ell = O(1)$) on one extreme and the brute force $2^{O(n)}$ on the other ($\ell = n^{1-o(1)}$). Our results show a similar behavior in the context of fatness and the packing problem: the running time of Theorem 1 is optimal, with the running time smoothly going from $2^{O(n^{1-1/d})}$ time in the case of $\alpha = O(1)$ to the trivial $2^{O(n)}$ time of brute force when $\alpha = n^{1/d}$.

Let $\mathcal{G}(d, L)$ denote the set of intersection graphs in \mathbb{R}^d where each object is an axis-parallel box whose side lengths form the multiset $\{1, \dots, 1, L\}$. Let us call such an axis-parallel box *canonical*. As usual, n denotes the number of objects (the number of vertices in the graph).

For example, it is easy to see that $1 \times 1 \times L$ boxes have stabbing number $O(L^{2/3})$. Any collection of $1 \times 1 \times L$ boxes of the same orientation can be stabbed by the lattice generated by the vertices of such a box, which has $O(L^2)$ points in a ball of radius $O(L)$. By taking the same lattice for the two other orientations, we obtain a complete stabbing set of size $O(L^2)$ inside a ball of radius $O(L)$ for all axis-parallel boxes of this shape. In general for $d \geq 3$, the stabbing number for canonical boxes is $\alpha = O(L^{1-1/d})$, so in particular, for $L = 1$ we have $\alpha = O(1)$, and for $L \geq n^{1/(d-1)}$ we have $\alpha = O(n^{1/d})$. In our main contribution, we show that this very restricted set of non-fat objects is sufficient to prove the desired lower bound.

► **Theorem 2.** *Let $d \geq 3$ be fixed. Then there is a constant $\gamma > 0$ such that for all $\alpha \in [1, n^{1/d}]$ it holds that INDEPENDENT SET on intersection graphs of d -dimensional canonical axis-parallel boxes of stabbing number α has no algorithm running in time $2^{\gamma n^{1-1/d} \alpha}$, unless ETH fails.*

An immediate corollary is that the $2^{O(n)}$ time brute-force algorithm cannot be improved, even for the intersection graph of axis-parallel boxes. This Corollary 3 can also be derived from a simpler construction by Chlebík and Chlebíková [7].

► **Corollary 3.** *Let $3 \leq d \in \mathbb{N}$ be fixed. Then INDEPENDENT SET on intersection graphs of axis-parallel boxes in d -dimensions has no algorithm running in time $2^{o(n)}$, unless ETH fails.*

In unit ball graphs, there is a lower bound of $2^{\Omega(n^{1-1/d})}$ under ETH, which of course carries over to intersection graphs of fat objects [9]. This latter reduction is based on establishing efficient routing constructions (called the “Cube Wiring theorem”) in the d -dimensional Euclidean grid. The crucial insight of the present paper is that tight lower bounds for *nonfat*

36:4 How Does Object Fatness Impact Packing?

objects can be obtained via INDEPENDENT SET on induced subgraphs of the d -dimensional *blown-up* grid cube, where each vertex is replaced by a clique of t vertices, fully connected to the adjacent cliques in all d directions. First we establish a lower bound for INDEPENDENT SET on subgraphs of such cubes (even for subgraphs of maximum degree 3), using and extending the Cube Wiring theorem [9]. Unlike for unit balls, it now seems difficult to realize every such subgraph G as intersection graph of appropriate boxes. Instead, we realize a graph G' that is obtained from G by some number of double subdivisions (subdividing some edge twice). As every double subdivision is known to increase the size of the maximum independent set by exactly 1, switching to G' does not cause a problem in the reduction.

The key insight of the reduction (in 3-dimensions) is that if $t = L^2$, then t vertices can be represented with $1 \times 1 \times L$ size boxes arranged in an $L \times L$ grid, occupying $O(L) \times O(L) \times O(L)$ space. Each t -clique of the blown-up cube is represented by such arrangements of boxes. The main challenge that we have to overcome is that the subgraph G may contain an arbitrary matching between two adjacent t -cliques. Given two sets of $1 \times 1 \times L$ size boxes arranged in two $L \times L$ grids, it seems unclear whether such arbitrary connections can be realized while staying in an $O(L) \times O(L) \times O(L)$ region of space. However, we show that this is possible, as the $L \times L$ grid arrangement allows easy reordering within the rows or within the columns, and it is known that any permutation of a grid can be obtained as doing a permutation first within the rows, then within the columns, and finally one more time within the rows. Thus with some effort, it is possible to build gadgets representing $L \times L$ vertices in an $O(L) \times O(L) \times O(L)$ region of space that allows arbitrary matchings to be realized with the adjacent gadgets.

The idea is similar in higher dimensions $d > 3$. We reduce from the INDEPENDENT SET problem on a subgraph of the blow-up of a d -dimensional grid where each vertex is blown-up into a clique of L^{d-1} vertices. Each gadget now contains L^{d-1} boxes of size $1 \times 1 \times \dots \times 1 \times L$ arranged in a grid. In order to implement arbitrary matchings between adjacent gadgets, we decompose every permutation of the $(d-1)$ -dimensional grid into $O(d)$ simpler permutations that are easy to realize in d -dimensional space.

We also study the complexity of packing in the context of parameterized algorithms: the question is how much one can improve the brute force $n^{O(k)}$ algorithm for finding k independent objects. We present a counterpart of Theorem 1 in this setting.

► **Theorem 4.** *Let $\alpha \in [1, \infty)$ and $2 \leq d \in \mathbb{N}$. There is a parameterized algorithm that solves independent set for intersection graphs of similarly sized α -stabbed objects in \mathbb{R}^d running in time $n^{O(k^{1-1/d}\alpha)}$, where the parameter k is the size of the maximum independent set.*

If one regards the parameterized algorithm's running time in terms of the instance size only, the result would be a $2^{O(n^{1-1/d}(\log n)^\alpha)}$ algorithm, which is slower than the running time $2^{O(n^{1-1/d}\alpha)}$ provided by the latter algorithm. The parameterized algorithm is based on a separator theorem by Miller *et al.* [27].

Finally, we sketch how the lower bound construction of Theorem 2 can be adapted to a parameterized setting, and obtain the following theorem:

► **Theorem 5.** *Let $3 \leq d \in \mathbb{N}$ be fixed. Then there is a constant $\gamma > 0$ such that for all $\alpha \in [1, n^{1/d}]$ it holds that deciding if there is an independent set of size k in intersection graphs of d -dimensional canonical axis-parallel boxes of stabbing number α has no $f(k)n^{\gamma k^{1-1/d}\alpha/\log k}$ algorithm for any computable function f , unless ETH fails.*

The crucial difference is that we are reducing from the PARTITIONED SUBGRAPH ISOMORPHISM problem instead of INDEPENDENT SET, which means that instead of choosing or not choosing a box (representing choosing or not choosing a vertex in the INDEPENDENT

SET problem), the solution needs to choose one of n very similar boxes (representing the choice of one of n vertices in a class of the partition). The overall structure of the reduction (e.g., routing in the blown-up d -dimensional grid) is similar to the proof of Theorem 2, and it can be found in the full version [23] along with other missing proofs.

2 The relationship between the stabbing number and fatness

In the usual definition of fatness, an object $o \subset \mathbb{R}^d$ is α -fat if there exists a ball of radius ρ_{in} contained in o and a ball of radius ρ_{out} that contains o , where $\rho_{\text{in}}/\rho_{\text{out}} = \alpha$. For a fixed constant α this is a useful definition and unifies many other similar notions in case of convex objects, i.e., it holds that a set of convex objects that is constant-fat for this notion of fatness are constant-fat for more restrictive definitions and vice versa. For our purposes however this definition is not fine-grained enough in the following sense. The fatness of a $1 \times 1 \times n$ box in three dimensions would be $\Theta(n)$, just as the fatness of a $1 \times n \times n$ box. As it will be apparent in what follows, we need a fatness definition according to which $1 \times n \times n$ boxes are much more fat than $1 \times 1 \times n$ boxes. For this purpose, we use the following weaker definition of fatness, that tracks the volume compared to a circumscribed ball more closely. (Note that constant-fat objects are also weakly constant-fat.)

► **Definition 6 (Weakly α -fat).** *A measurable object $o \subseteq \mathbb{R}^d$ is α -fat for some $\alpha \in [1, \infty)$ if $\text{Vol}(o)/\text{Vol}(B) \leq \alpha^d$, where $\text{Vol}(o)$ and $\text{Vol}(B)$ denotes the volume of o and the volume of its circumscribed ball B respectively.*

An object o is *strongly α -fat* if for any ball B centered inside o we have $\text{Vol}(B \cap o)/\text{Vol}(B) \geq \alpha^d$. In case of convex objects, weak fatness coincides with strong fatness up to constant factors, see [31].

The next theorem shows that the inverse of the weak fatness of an object family is related to the stabbing number. In a sense, this means that the stabbing number is a further weakening of weak fatness. Note that in our setting, the stabbing number will be polynomial in n (i.e., $\alpha = n^\lambda$ for some constant λ), so the $\log n$ term is insignificant.

► **Theorem 7.** *Let d be a fixed constant. Then the stabbing number of any family of n weakly $(1/\alpha)$ -fat (measurable) objects in \mathbb{R}^d is $O(\alpha \log^{1/d} n)$.*

Proof. Consider a family F of weakly $1/\alpha$ -fat objects. Let B be a ball of radius δ , and let F_B be the set of objects contained in B of diameter at least $\delta/2$. It is sufficient to show that we can stab F_B with $O(\alpha^d \log n)$ points. Pick $k = \lfloor (4\alpha)^d (\log n + 1) \rfloor$ points p_1, \dots, p_k independently uniformly at random in B . For any given object o , its volume is at least $\text{Vol}(B)/(4\alpha)^d$, so the probability that a given p_i is not in o is at most $1 - 1/(4\alpha)^d$. Since the k points are chosen independently, the probability that a given object o is unstabbed is at most $\left(1 - \frac{1}{(4\alpha)^d}\right)^k$. By the union bound, the probability that there is an unstabbed object is at most

$$n \left(1 - \frac{1}{(4\alpha)^d}\right)^k = n \left(1 - \frac{1}{(4\alpha)^d}\right)^{\lfloor (4\alpha)^d (\log n + 1) \rfloor} < n(1/e)^{\log n + 1} < 1.$$

Consequently, there exists an outcome where all objects are stabbed. ◀

We conclude this section with the following theorem, which shows an even stronger connection between fatness and stabbing in case of convex objects. The theorem uses the existence of the John ellipsoid [22] and the ε -net theorem [20].

36:6 How Does Object Fatness Impact Packing?

► **Theorem 8.** *Let d be a fixed constant. Then the stabbing number of any family of n weakly $(1/\alpha)$ -fat convex objects in \mathbb{R}^d is $O(\alpha \log^{1/d} \alpha)$.*

Proof. Consider a family F of weakly $1/\alpha$ -fat convex objects. Let B be a ball of radius δ , and let F_B be the set of objects contained in B of diameter at least $\delta/2$. It is sufficient to show that we can stab F_B with $O(\alpha^d \log \alpha)$ points. For any given object o , its volume is at least $\text{Vol}(B)/(4\alpha)^d$. Every convex object $o \in F_B$ contains an ellipsoid $\ell(o) \subseteq o$ such that $\text{Vol}(o)/\text{Vol}(\ell(o)) \leq d^d$ [22]. Since the VC-dimension of ellipsoids in \mathbb{R}^d is $O(d^2)$ [2], the ϵ -net theorem [20] implies that the ellipsoids $\ell(o)$ ($o \in F_B$) can be stabbed by $O(\frac{d^2}{1/(4\alpha)^d} \log \frac{d^2}{1/(4\alpha)^d}) = O(\alpha^d \log \alpha)$ points. Since the ellipsoids are contained in their respective objects, this point set also stabs all objects in F_B . ◀

3 Algorithms

We require very little from the objects that we use in our algorithms. It is necessary that we can decide in polynomial time whether a point is contained in an object, whether two objects intersect, and whether an object intersects some given sphere, ball, and empty or dense hypercube. Let us assume that such operations are possible from now on.

To prove Theorem 4, we use the following separator theorem, due to Miller et al. [27]. The *ply* of a set of objects in \mathbb{R}^d is the largest number p such that there exists a point $x \in \mathbb{R}^d$ which is contained in p objects.

► **Theorem 9** (Miller et al. [27]). *Let $\Gamma = \{B_1, \dots, B_n\}$ be a collection of n closed balls in \mathbb{R}^d with ply at most p . Then there exists a sphere S whose boundary intersects at most $O(p^{1/d} n^{1-1/d})$ balls, and the number of balls in Γ disjoint from S that fall inside and outside S are both at most $\frac{d+1}{d+2}n$.*

Proof sketch. Consider the set of balls B made up by the circumscribed balls of the objects in a maximum independent set. We claim that the ply of this set is $O(\alpha^d)$. To prove the claim, let S be a subset of the independent set whose circumscribed balls overlap at a point $x \in \mathbb{R}^d$. Since the objects are similarly sized, S must lie within a ball centered at x whose radius is at most a constant times the diameter of the largest object. Thus, S can be stabbed by $O(\alpha^d)$ points. However, as S forms an independent set, each point can only stab at most one object from S . Therefore, $|S| = O(\alpha^d)$. By Theorem 9 the ball set B has a sphere separator intersecting $O((\alpha^d)^{1/d} k^{1-1/d}) = O(k^{1-1/d} \alpha)$ balls. We proceed by guessing such a sphere: a discretization argument shows that there are $\text{poly}(n)$ distinct guesses for this sphere separator. We also guess the set of objects in the independent set that intersect the sphere, and remove all other objects intersecting the sphere or the guessed objects, and recurse on the remaining objects inside S and on the remaining objects outside S . The resulting running time is $n^{O(k^{1-1/d} \alpha)}$. ◀

For arbitrary size objects that are $O(1)$ -fat in some stronger sense (or just $O(1)$ -stabbed), we can apply the above scheme of guessing a separating sphere or hypercube, and use one of the many separator theorems designed for objects of small ply. See [6, 19, 29]. One can also apply [9] since in case of ply 1, the weights are constants; although the theorem is stated for the usual notion of fatness, the proof itself uses only the stabbing number. We get the following theorem.

► **Theorem 10.** *Let $2 \leq d \in \mathbb{N}$. There is a parameterized algorithm that solves INDEPENDENT SET for intersection graphs of $O(1)$ -stabbed objects in \mathbb{R}^d running in time $n^{O(k^{1-1/d})}$, where the parameter k is the size of the maximum independent set.*

The algorithm for Theorem 1 is an adaptation of the INDEPENDENT SET algorithm for fat objects from [9], based on weighted cliques, and its proof is deferred to the full version [23].

4 Wiring in a blowup of the Euclidean Cube

Our wiring theorem relies on the folklore observation that can be informally stated the following way: an $n \times m$ matrix can be sorted by first permuting the elements within each row, then permuting the elements within each column, and then permuting the elements in each row again. Note that the permutations are independent of each other, and they are not sorting steps; the permutations required are quite specialized. We state the lemma in a more group-theoretic setting. Let $\text{Sym}(X)$ denote the symmetric group on the set X .

► **Lemma 11** (Lemma 4 of [1]). *Let A and B be two finite sets. Then $\text{Sym}(A \times B) = G_A G_B G_A$, where G_A is the subgroup of $\text{Sym}(A \times B)$ consisting of permutations π where $\pi(a, b) \in A \times \{b\}$ for all $(a, b) \in A \times B$, and G_B is the subgroup of $\text{Sym}(A \times B)$ consisting of permutations π where $\pi(a, b) \in \{a\} \times B$ for all $(a, b) \in A \times B$.*

► **Corollary 12.** *Let $2 \leq d \in \mathbb{N}$ and let A_1, A_2, \dots, A_d be finite sets. Then $\Gamma \stackrel{\text{def}}{=} \text{Sym}(A_1 \times A_2 \times \dots \times A_d)$ is of the form $\Gamma = G_1 G_2 \dots G_{d-1} G_d G_{d-1} G_{d-2} \dots G_1$, where G_i is the subgroup of Γ consisting of permutations π where $\pi(a_1, \dots, a_i, \dots, a_d) \in \{a_1\} \times \dots \times \{a_{i-1}\} \times A_i \times \{a_{i+1}\} \times \dots \times \{a_d\}$ for all $(a_1, \dots, a_d) \in \Gamma$.*

Proof. We use induction on d ; for $d = 2$, the statement is equivalent to Lemma 11. Let $d \geq 3$. We can write Γ as $\text{Sym}((A_1 \times \dots \times A_{d-1}) \times A_d)$, so by induction (for $d = 2$), we have that $\Gamma = G_1 \times G_{A_2 \times \dots \times A_d} \times G_1$. By induction, we also have that $G_{A_2 \times \dots \times A_d} = G_2 \dots G_{d-1} G_d G_{d-1} G_{d-2} \dots G_2$, therefore $\Gamma = G_1 G_2 \dots G_{d-1} G_d G_{d-1} G_{d-2} \dots G_1$. ◀

For an integer n , let $[n] = \{1, \dots, n\}$. Let $\mathcal{EC}^d(n)$ be the d -dimensional Euclidean grid graph whose vertices are $[n]^d$, and $x, y \in V(G)$ are connected if and only if they are at distance 1 in \mathbb{R}^d . For $x \in \mathbb{Z}^d$ and $S \subset \mathbb{Z}^d$, we use the shorthand $x + S \stackrel{\text{def}}{=} \{x + y \mid y \in S\}$. Let $\mathcal{BEC}^d(n, t)$ denote the t -fold blowup of $\mathcal{EC}^d(n)$, where all vertices of $\mathcal{EC}^d(n)$ are exchanged with a clique of size t , and vertices in neighboring cliques are connected. More precisely,

$$\begin{aligned} V(\mathcal{BEC}^d(n, t)) &= [n]^d \times [t] \\ E(\mathcal{BEC}^d(n, t)) &= \{(x, i)(y, j) \mid x = y \vee (x, y) \in E(\mathcal{EC}^d(n))\}. \end{aligned}$$

Our second key ingredient is the Euclidean Cube Wiring theorem.

► **Theorem 13** (Theorem 21 in [9]). *Let $3 \leq d \in \mathbb{Z}$. There exists a constant c dependent only on the dimension such that any matching M between $P = [n]^{d-1} \times \{1\}$ and $Q \stackrel{\text{def}}{=} [n]^{d-1} \times \{cn\}$ can be embedded in $\mathcal{EC}^d(cn)$, that is, there is a set of vertex disjoint paths connecting p and q in $\mathcal{EC}^d(cn)$ for all $pq \in M$.*

► **Theorem 14** (Blown-up Cube Wiring). *Let $3 \leq d \in \mathbb{Z}$, and let n, t be positive integers. We consider two opposing facets of the blown-up cube $\mathcal{C} \stackrel{\text{def}}{=} \mathcal{BEC}^d(cn, t)$ (where $c \in \mathbb{Z}_+$ depends only on d):*

$$\begin{aligned} P &\stackrel{\text{def}}{=} ([n]^{d-1} \times \{1\}) \times [t] \\ Q &\stackrel{\text{def}}{=} ([n]^{d-1} \times \{cn\}) \times [t] \end{aligned}$$

Any matching M between P and Q can be embedded in \mathcal{C} , that is, there is a constant integer c dependent only on the dimension d such that for any matching M there is a set of vertex disjoint paths connecting p and q in $\mathcal{BEC}^d(cn, t)$ for all $pq \in M$.

Proof. Without loss of generality, suppose that M is a perfect matching between P and Q (this can be ensured by adding dummy edges to M if necessary). Let $c = c' + 2$ where c' is a constant such that cube wiring can be done in height $h = c'n$. Let $A = [n]^{d-1}$ and let $B = [t]$. The matching M can be regarded as a permutation π of $A \times B$, where $\pi(a, b) = (a', b')$ if $((a, b)(a', b')) \in M$.

By Lemma 11, there exists a permutation $\pi_A \in G_A$ and $\pi_B, \pi'_B \in G_B$ such that $\pi = \pi'_B \pi_A \pi_B$, where G_A and G_B are defined as in Lemma 11. We can think of both π_B and π'_B as the union of n^{d-1} distinct permutations of $[t]$. We can realize π_B using one matching: for all $(x, i) \in A \times B$, we add the edge $((x, 1), i)((x, 2), j)$ to M_B , where $\pi_B(x, i) = (x, j)$. As a result, M_B is a perfect matching between P and the next layer of the blown-up cube, $P' \stackrel{\text{def}}{=} ([n]^{d-1} \times \{2\}) \times [t]$. Similarly, for all $(x, i) \in A \times B$, let M'_B contain the edge $((x, cn - 1), i)((x, cn), j)$, where $\pi'_B(x, i) = (x, j)$; this matches $Q' \stackrel{\text{def}}{=} ([n]^{d-1} \times \{cn - 1\}) \times [t]$ to Q . Finally, by the Cube Wiring Theorem (Theorem 13), for each $i \in [t]$, there are vertex disjoint paths from $P'_i \stackrel{\text{def}}{=} ([n]^{d-1} \times \{2\}) \times \{i\}$ to $Q'_i \stackrel{\text{def}}{=} ([n]^{d-1} \times \{cn - 1\}) \times \{i\}$ that realizes the matching

$$M_A^i \stackrel{\text{def}}{=} \{(x, i)(y, i) \mid x \in [n]^{d-1} \text{ and } \pi_A(x, i) = (y, i)\}.$$

For each $i \in [t]$, these wirings are vertex disjoint since they are contained in vertex disjoint Euclidean grid hypercubes. The matchings M_A^i for $i \in [t]$ together with the matchings M_B and M'_B realize the matching M . ◀

5 Lower bounds for packing isometric axis-parallel boxes

Our first lower bound shows that the running time of the algorithm in Theorem 1 is tight under ETH.

Overview of the proof of Theorem 2. Our proof is a reduction from (3,3)-SAT, the satisfiability problem of CNF formulas where clauses have size at most three and each variable occurs at most three times. Such formulas have the property that if they have n variables, then they have $O(n)$ clauses. The problem has no $2^{o(n)}$ algorithm under ETH [10].

The proof has two steps; the first step is a reduction from (3,3)-SAT to INDEPENDENT SET in certain subgraphs of the blown-up Euclidean cube, and the second step is to show that these subgraphs can essentially be realized with axis-parallel boxes. Throughout the proof, we consider the dimension d to be a constant.


The *incidence graph* of a (3,3)-CNF formula ϕ is a graph where vertices correspond to clauses and variables of ϕ , and a variable and clause vertex are connected if and only if the variable occurs in the clause.

5.1 Independent Set in subgraphs of the blown-up Euclidean cube

A simple and generic lower bound construction for Independent Set

We give a generic reduction from (3,3)-SAT to INDEPENDENT SET, which serves as a skeleton for the more geometric type of reduction we will do later.

Consider the incidence graph of ϕ . Replace each variable vertex v with a cycle of length 6, consisting of vertices v^1, \dots, v^6 , where the edges formerly incident to v are now connected to distinct cycle vertices v^2, v^4 or v^6 for positive literals and to v^1, v^3 or v^5 for negative literals (see Figure 1). We replace each clause vertex w that corresponds to a clause of exactly 3 literals with a cycle of length three, and connect the formerly incident edges to distinct



■ **Figure 1** The graph G_ϕ for $\phi = (v_1 \vee \neg v_2 \vee v_3)$.

vertices of the triangle. For clauses that have exactly two literals, the gadget is a single edge, and we connect the formerly incident edges to distinct endpoints of the edge. We can eliminate clauses of size 1 in a preprocessing step. Let G'_ϕ be the resulting graph.

An independent set can contain at most 3 vertices of a variable cycle of length 6, and at most 1 vertex per clause gadget. Observe that a formula with ν variables and γ clauses has an independent set of size $3\nu + \gamma$ if and only if the original formula is satisfiable.

Let G be a graph and let uv be an edge of G . A *double subdivision* of uv is replacing uv with a path of length 3, i.e., we add the new vertices w and w' , remove the edge uv and add the edges $uw, ww', w'v$. A graph that can be obtained from G by some sequence of double subdivisions is called an *even subdivision* of G . Observe that a double subdivision increases the size of the maximum independent set by one, so G has an independent set of size k if and only if its even subdivision G' has an independent set of size $k + \frac{|V(G')| - |V(G)|}{2}$.

Embedding G'_ϕ into a blown-up cube

In a blown-up cube $\mathcal{BEC}^d(n, t)$, we call a clique corresponding to $x \in [n]^d$ the *cell* of x or simply a cell, that is, the cell of x is defined as the set of vertices $\{x\} \times [t] \subset V(\mathcal{BEC}^d(n, t))$.

The following is a tight lower bound for INDEPENDENT SET inside the blown-up Euclidean cube.


► **Theorem 15.** *For any fixed constant $d \geq 3$, there exists a $\gamma > 0$ such that for any $t \geq 2$ there is no $2^{\gamma n^{1-1/d} t^{1/d}}$ algorithm for INDEPENDENT SET for subgraphs of the blown-up cube $\mathcal{C} \stackrel{\text{def}}{=} \mathcal{BEC}^d((n/t)^{1/d}, t)$ under ETH. The lower bound holds even if the subgraph G has maximum degree three, and the neighbors of each vertex in G lie in distinct cells.*

Proof sketch. Given a (3, 3)-SAT formula ϕ , we show that we construct a subgraph of a blown-up cube with the required properties that is also an even subdivision of G'_ϕ . If ϕ has \bar{n} literals, then we create a subgraph G that has $n = c \cdot \bar{n}^{\frac{d}{d-1}} / t^{\frac{1}{d-1}}$ vertices; a simple computation shows that this is sufficient. The variable cycles become cycles of length 6, and they are placed densely within cells that lie in some facet of \mathcal{C} . Similarly, for clauses of size two and three, we associate an edge or a triangle in the cells of the opposing facet of \mathcal{C} . Using Theorem 14, we can construct wires that for each literal connect the relevant vertex of the variable cycle to the relevant vertex of the clause cycle. If the resulting wire has even length, then we add an extra edge to its end that connects to the clause cycle. The resulting embedding has the desired properties. ◀

5.2 Detailed construction and gadgetry

Having established our lower bound for blown-up Euclidean cubes, we now need to construct a set of canonical boxes whose intersection graph is an even subdivision of a given subgraph with maximum degree three where the neighbors of each vertex lie in distinct cells.

36:10 How Does Object Fatness Impact Packing?



■ **Figure 2** A basic brick.

► **Theorem 16.** *Let $d \geq 3$ and $L \geq 16$ be fixed, and let G be a subgraph of the blown-up cube $\mathcal{C} = \mathcal{BEC}^d(s, (L/8)^{d-1})$ of maximum degree three, where the neighbors of each vertex lie in distinct cells. Then G has an even subdivision G' that can be realized using boxes of size $1 \times \dots \times 1 \times L$. Moreover, given G , the boxes of G' can be constructed in $O(|V(\mathcal{C})|)$ time, and $|V(G')| = O(|V(G)|)$.*

We consider $d = 3$ first; later on, we show how the construction can be generalized to higher dimensions. We need to define a set of boxes whose intersection graph is an even subdivision of G . The idea is to create a generic *module* that is able to represent a subgraph of G induced by any cell; these modules will take up $O(L) \times O(L) \times O(L)$ space. The modules are arranged into a larger cube of side length $O(sL)$ to make up the final construction.


Due to space constraints, we concentrate on giving a picture of the overall construction for $d = 3$, and on presenting our most intricate gadget that is capable of realizing so-called parallel matchings. The rest of the gadgetry and other details are deferred to Appendix A.

Modules and bricks

We index the vertices in a cell by a pair from $[L/8]^2$. The starting object in our reduction is a set of $(L/8)^2$ disjoint boxes parallel to the same axis, arranged loosely in an $L/8 \times L/8$ grid structure called a *brick*. See Figure 2 for an example. Loosely speaking, each box of each brick within the cell's module can be associated with a vertex of the cell; for a brick B , we can refer to a box corresponding to vertex (i, j) of the cell as $B(i, j)$.

Let X be the set of cells within \mathcal{C} : $X \stackrel{\text{def}}{=} \{\{x\} \times [L/8]^2 \mid x \in [cn]^d\}$. The wiring within each cell $x \in X$ will be represented by $O(1)$ bricks, and these bricks will fit in an $O(L)$ side length module.

The position of a brick can be specified by defining its axis (along which the side length of the boxes is L), and for each box (i, j) within the brick, defining the coordinates of its lexicographically smallest corner (or *lexmin corner* for short). For example, consider the brick B with axis x_3 where box $B(i, j)$ has coordinates $(3i, 3j, 0)$. (See Figure 2.) This brick and all bricks isometric to this are called *basic bricks*. Most bricks can be thought of as a perturbation of a basic brick, where we apply shifts to each box. The eventual module that we create will consist of several bricks, which together will represent an even subdivision of the sparse graph G restricted to a given cell. Note that no single brick can be said to represent the set of vertices in a cell. When defining our gadgetry, it is convenient to talk about these bricks, even though in the final construction we only need a certain subset of the boxes within each brick. We can remove the unwanted boxes from each brick at a later stage.



■ **Figure 3** Left: First column of a parallel matching gadget for the permutation $\pi_1(1) = 1, \pi_1(2) = 4, \pi_1(3) = 2, \pi_1(4) = 3$. Boxes of each color induce paths; boxes of different color are disjoint. Right: A full parallel matching gadget.

The parallel matching gadget

A parallel matching gadget is capable of realizing a matching between two cells where the endpoints of each matching edge differ only on a fixed coordinate, so for $d = 3$, all edges are of the type $((x, (i, j)), (x', (i', j)))$ or all edges are of the type $((x, (i, j)), (x', (i, j')))$ for some cells x and x' . We call a matching with this property a *parallel matching*. Parallel matchings can be decomposed into matchings on disjoint cliques, where each clique contains vertices that share all of their coordinates except one. In the remainder of this gadget's description, we will omit the cells x and x' from the coordinate lists.

Suppose that each matching edge is of the form $((i, j), (i', j))$. Let $\pi_j(i)$ denote the first coordinate of the pair of (i, j) , that is, suppose that the matching edges are $((i, j), (\pi_j(i), j))$, $i \in I_j$ for some sets $I_j \subseteq [L/8]$. Instead of realizing these matchings, we first extend them to permutations π_j on each clique $[L/8] \times \{j\}$. A permutation can be thought of as a perfect matching between two copies of a set; by removing the unwanted vertices (removing the unwanted boxes) we can get to a representation of the matching, i.e., a set of vertex disjoint paths that connect box (i, j) in the starting brick to box $(\pi_j(i), j)$ in the target brick.

In every brick, each box is translated individually, where the translation vector's component along the brick's axis must be an integer $k \in 3 \cdot \{-L/8, \dots, L/8\}$, and along the other axes it must be of the form k/L for some $k \in \{-L/8, \dots, L/8\}$. For a brick B , its box of index (i, j) is denoted by $B(i, j)$, and recall that the position of a box is defined by its lexmin corner and the axis of the brick.

We give the coordinates of each box in each brick of the parallel matching gadget below. Let us take the matching edges where $j = 1$ first. We start with the first column of the brick ($j = 1$), where the coordinates of $B^{(1)}(i, 1)$ are $(3i, 3 + i/L, -3i)$. See the left hand side of Figure 3 that illustrates the idea behind the gadget. The coordinates for $B^{(1)}(i, j)$ are $(3i, 3j + i/L, -3i)$. The first column of the next brick $B^{(2)}$ has axis x_1 and the coordinates of $B^{(2)}(i, 1)$ are $(0, 4 + i/L, L - 1 - 3i)$, that is, these boxes touch the previously defined boxes of $B^{(1)}$ from "behind" in Figure 3. In general, $B^{(2)}(i, j)$ has coordinates $(0, 3j + 1 + i/L, L - 3i)$. The next brick $B^{(3)}$ also has axis x_1 , and the coordinates for $B^{(3)}(i, j)$ are $(L/2 + 3\pi_j(i), 3j + 1 + \pi_j(i)/L, L - 3i)$, that is, we change the box perturbations along the first and second coordinate. Finally, the last brick $B^{(4)}$ has axis x_3 and the coordinates are $(3L/2 + 3\pi_j(i), 3j - \pi_j(i)/L, L - 3i)$, i.e., they are placed "in front of"

36:12 How Does Object Fatness Impact Packing?

the bricks of $B^{(3)}$ in Figure 3. This can be rewritten as $B^{(4)}(i', j)$ having coordinates $(3L/2 + 3i', 3j - i'/L, L - 3\pi_j^{-1}(i'))$. Notice that in the final brick, we indeed have the desired ordering, i.e., the ordering of the boxes along the x_1 axis is as required. It is routine to check that the intersection graph induced each column of this parallel matching gadget consists of vertex disjoint paths of length four. Different columns are also disjoint since projecting the boxes of column j onto the x_2 axis results in a subset of the open interval $(3j - 0.5, 3j + 2.5)$.

Using several parallel matching gadgets, by Lemma 11 we are capable of representing arbitrary matchings between two neighboring cells or within a single cell in $O(L) \times O(L) \times O(L)$ space. Further detailed gadgetry describing how branching gadgets are made (capable of representing a collection of degree 3 vertices), and how everything can be fit into modules of side length $O(L)$ are described in Appendix A. Using Theorem 16, it is easy to prove Theorem 2.

Proof of Theorem 2. Set $L \stackrel{\text{def}}{=} \max(16, \alpha^{\frac{d}{d-1}})$. This choice of L implies that any family of canonical boxes of size $1 \times 1 \times L$ are $O(\alpha)$ -stabbed. Furthermore, set $t = (L/8)^{d-1}$. The proof is by reduction from INDEPENDENT SET on subgraphs of the blown-up cube $\mathcal{C} \stackrel{\text{def}}{=} \mathcal{BEC}^d((\bar{n}/t)^{1/d}, t)$, where the subgraph G has maximum degree three, and the neighbors of each vertex in G lie in distinct cells. By Theorem 15, there is no $\gamma > 0$ for which a $2^{\gamma n^{1-1/d} t^{1/d}}$ algorithm exists for this problem under ETH.

Let G be a subgraph of \mathcal{C} as described above. By Theorem 16, we can realize an odd subdivision G' of G using boxes of size $1 \times \dots \times 1 \times L$, with $O(\bar{n})$ vertices in $\text{poly}(\bar{n})$ time. If for any $\gamma > 0$ there is an algorithm for INDEPENDENT SET on α -stabbed canonical boxes with running time $2^{\gamma n^{1-1/d} \alpha}$, then this translates into $2^{\gamma n^{1-1/d} L^{1-1/d}}$ algorithms for all $\gamma > 0$. This can be composed with our construction to get $2^{\gamma \bar{n}^{(1-1/d)} t^{1/d}}$ algorithms for all $\gamma > 0$ for INDEPENDENT SET on the described subgraphs of \mathcal{C} , which contradicts ETH according to Theorem 15. \blacktriangleleft

6 Conclusion

We have explored the impact of the stabbing number on the complexity of packing. We have seen that subexponential packing algorithms are possible for similarly sized objects if the stabbing number is $o(n^{1/d})$. The subexponential algorithms could be derived from powerful separator theorems, while the lower bounds required custom wiring results and non-trivial geometric gadgetry. We propose two open problems for future research.

- What is the precise impact of the stabbing number on the complexity of packing if objects are not similarly sized? One can get a subexponential algorithm by an adaptation of the separator in [9], but it yields an algorithm whose dependence on α is much weaker: it has α^d in the exponent instead of α . Is this algorithm optimal?
- Is there a subexponential algorithm for the DOMINATING SET problem in intersection graphs of α -stabbed similarly sized objects? Or even for n axis-parallel $1 \times n^\epsilon$ and $n^\epsilon \times 1$ boxes in two dimensions?

References

- 1 Miklós Abért. Symmetric groups as products of abelian subgroups. *Bulletin of the London Mathematical Society*, 34(4):451–456, 2002.
- 2 Yohji Akama and Kei Irie. VC dimension of ellipsoids. *CoRR*, abs/1109.4347, 2011. [arXiv:1109.4347](https://arxiv.org/abs/1109.4347).

- 3 Jochen Alber and Jirí Fiala. Geometric separation and exact solutions for the parameterized independent set problem on disk graphs. *Journal of Algorithms*, 52(2):134–151, 2004. doi:10.1016/j.jalgor.2003.10.001.
- 4 Julien Baste and Dimitrios M. Thilikos. Contraction-Bidimensionality of Geometric Intersection Graphs. In *IPEC 2017*, volume 89 of *LIPICs*, pages 5:1–5:13, 2018. doi:10.4230/LIPICs.IPEC.2017.5.
- 5 Csaba Biró, Édouard Bonnet, Dániel Marx, Tillmann Miltzow, and Paweł Rzażewski. Fine-grained complexity of coloring unit disks and balls. *JoCG*, 9(2):47–80, 2018. doi:10.20382/jocg.v9i2a4.
- 6 Timothy M. Chan. Polynomial-time approximation schemes for packing and piercing fat objects. *J. Algorithms*, 46(2):178–189, 2003. doi:10.1016/S0196-6774(02)00294-8.
- 7 Miroslav Chlebík and Janka Chlebíková. Approximation hardness of optimization problems in intersection graphs of d -dimensional boxes. In *Proceedings of SODA 2005*, pages 267–276. SIAM, 2005. URL: <http://dl.acm.org/citation.cfm?id=1070432.1070470>.
- 8 Marek Cygan, Fedor V Fomin, Łukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 9 Mark de Berg, Hans L. Bodlaender, Sándor Kisfaludi-Bak, Dániel Marx, and Tom C. van der Zanden. A framework for ETH-tight algorithms and lower bounds in geometric intersection graphs. In *Proceedings of STOC 2018*, pages 574–586, 2018. doi:10.1145/3188745.3188854.
- 10 Mark de Berg, Hans L. Bodlaender, Sándor Kisfaludi-Bak, Dániel Marx, and Tom C. van der Zanden. A Framework for ETH-Tight Algorithms and Lower Bounds in Geometric Intersection Graphs. *CoRR*, abs/1803.10633, 2018. arXiv:1803.10633.
- 11 Erik D. Demaine, Fedor V. Fomin, Mohammad Taghi Hajiaghayi, and Dimitrios M. Thilikos. Fixed-parameter algorithms for (k, r) -Center in planar graphs and map graphs. *ACM Transactions on Algorithms*, 1(1):33–47, 2005.
- 12 Erik D. Demaine, Fedor V. Fomin, Mohammad Taghi Hajiaghayi, and Dimitrios M. Thilikos. Subexponential parameterized algorithms on bounded-genus graphs and H -minor-free graphs. *Journal of the ACM*, 52(6):866–893, 2005. doi:10.1145/1101821.1101823.
- 13 Frederic Dorn, Fedor V. Fomin, and Dimitrios M. Thilikos. Subexponential parameterized algorithms. *Computer Science Review*, 2(1):29–39, 2008.
- 14 Frederic Dorn, Fedor V. Fomin, and Dimitrios M. Thilikos. Catalan structures and dynamic programming in H -minor-free graphs. *J. Comput. Syst. Sci.*, 78(5):1606–1622, 2012.
- 15 Frederic Dorn, Eelko Penninx, Hans L. Bodlaender, and Fedor V. Fomin. Efficient Exact Algorithms on Planar Graphs: Exploiting Sphere Cut Decompositions. *Algorithmica*, 58(3):790–810, 2010.
- 16 Fedor V. Fomin, Daniel Lokshtanov, Venkatesh Raman, and Saket Saurabh. Subexponential algorithms for partial cover problems. *Inf. Process. Lett.*, 111(16):814–818, 2011.
- 17 Fedor V. Fomin, Daniel Lokshtanov, and Saket Saurabh. Bidimensionality and geometric graphs. In *Proceedings of SODA 2012*, pages 1563–1575. SIAM, 2012. URL: <http://portal.acm.org/citation.cfm?id=2095240&CFID=63838676&CFTOKEN=79617016>.
- 18 Fedor V. Fomin and Dimitrios M. Thilikos. Dominating Sets in Planar Graphs: Branch-Width and Exponential Speed-Up. *SIAM J. Comput.*, 36(2):281–309, 2006.
- 19 Sarel Har-Peled and Kent Quanrud. Approximation Algorithms for Polynomial-Expansion and Low-Density Graphs. *SIAM J. Comput.*, 46(6):1712–1744, 2017. doi:10.1137/16M1079336.
- 20 David Haussler and Emo Welzl. ε -nets and simplex range queries. *Discrete & Computational Geometry*, 2(2):127–151, June 1987. doi:10.1007/BF02187876.
- 21 Russell Impagliazzo and Ramamohan Paturi. On the Complexity of k -SAT. *Journal of Computer and System Sciences*, 62(2):367–375, 2001. doi:10.1006/jcss.2000.1727.
- 22 Fritz John. *Extremum Problems with Inequalities as Subsidiary Conditions*, pages 197–215. Springer Basel, Basel, 2014. doi:10.1007/978-3-0348-0439-4_9.

36:14 How Does Object Fatness Impact Packing?

- 23 Sándor Kisfaludi-Bak, Dániel Marx, and Tom C. van der Zanden. How does object fatness impact the complexity of packing in d dimensions? *CoRR*, abs/1909.12044, September 2019. [arXiv:1909.12044](https://arxiv.org/abs/1909.12044).
- 24 Philip N. Klein and Dániel Marx. A subexponential parameterized algorithm for Subset TSP on planar graphs. In *SODA 2014 Proc.*, pages 1812–1830, 2014.
- 25 Dániel Marx and Michal Pilipczuk. Optimal Parameterized Algorithms for Planar Facility Location Problems Using Voronoi Diagrams. In *Proceedings of ESA 2015*, volume 9294 of *LNCS*, pages 865–877. Springer, 2015. doi:10.1007/978-3-662-48350-3_72.
- 26 Dániel Marx and Anastasios Sidiropoulos. The limited blessing of low dimensionality: when $1 - 1/d$ is the best possible exponent for d -dimensional geometric problems. In *Proceedings of SoCG 2014*, pages 67–76. ACM, 2014. doi:10.1145/2582112.2582124.
- 27 Gary L. Miller, Shang-Hua Teng, William P. Thurston, and Stephen A. Vavasis. Separators for sphere-packings and nearest neighbor graphs. *J. ACM*, 44(1):1–29, 1997. doi:10.1145/256292.256294.
- 28 Marcin Pilipczuk, Michał Pilipczuk, Piotr Sankowski, and Erik Jan van Leeuwen. Subexponential-Time Parameterized Algorithm for Steiner Tree on Planar Graphs. In *STACS 2013 Proc.*, pages 353–364, 2013.
- 29 Warren D. Smith and Nicholas C. Wormald. Geometric Separator Theorems & Applications. In *Proceedings of the 39th Annual Symposium on Foundations of Computer Science, FOCS 1998*, pages 232–243. IEEE Computer Society, 1998. doi:10.1109/SFCS.1998.743449.
- 30 Dimitrios M. Thilikos. Fast Sub-exponential Algorithms and Compactness in Planar Graphs. In *ESA 2011 Proc.*, pages 358–369, 2011.
- 31 A. Frank van der Stappen, Dan Halperin, and Mark H. Overmars. The Complexity of the Free Space for a Robot Moving Amidst Fat Obstacles. *Comput. Geom.*, 3:353–373, 1993. doi:10.1016/0925-7721(93)90007-S.

A Gadgetry and further construction details for the proof of Theorem 16

Parity Fix, adjustment, bridge, and elbow gadgets


The parity fix gadget is introduced so that we can ensure that each of the subdivisions that we create are even subdivisions. The gadget induces a path of length 3 or 4 depending on our needs, but occupies the same space in both cases. More precisely, the parity fix gadget contains three or four boxes, depending on the parity we need. The union of the boxes is a larger box of size $3L \times 1 \times 1$; it is easy to see that within that space we can realize both a path of length three and four using $L \times 1 \times 1$ boxes: one can cover the larger box by placing their lexmin corners at equal length intervals.

We can bridge distance along the axis of a basic brick by putting basic bricks next to each other, where each box intersects only the box of the same index from the previous and following brick. This creates a set of $(L/8)^2$ vertex disjoint paths in the intersection graph. We call this a *bridge gadget*.

Using two bricks of the same axis, we can in one step get rid of a perturbation (or introduce one). Let B be a normal brick with axis x_3 that is a perturbation of the basic brick. We introduce the basic brick B' that is the translate of the basic brick with the vector $(0, 1, L/2)$. Notice that box $B(i, j)$ intersects $B'(i, j)$ and no other boxes. Moreover, we could even introduce arbitrary perturbations along the x_1 axis in B' and along the x_3 axis within both B and B' without changing the intersection graph induced by B and B' . We call a pair of normal bricks that are a translated and rotated version of these an *adjustment gadget*.



■ **Figure 4** An elbow.




■ **Figure 5** The first “column” of a branching gadget.

Next, we introduce a way to change brick axis using an “elbow”. Consider a brick B that is a perturbation of the basic brick, where box (i, j) has coordinates $(3i, 3j, -3i)$. The brick B' has axis x_1 and the coordinates for $B'(i, j)$ are $(3i, 3j, L - 3i)$ (see Figure 4). Notice that using these *elbow gadgets* and adjustment gadgets together, one can route from any brick to any other brick at distance $\Theta(L)$ in $O(1)$ steps.

Realizing an arbitrary matching of a biclique or clique

We can regard a general matching M induced by two neighboring cells as a permutation of $[L/8]^2$, which can be written as the product of three special permutations by Corollary 12 that correspond to parallel matchings; i.e., the matching M is realizable as the succession of three parallel matchings. This means that each edge of M becomes a path of length three, so by using three parallel matching gadgets in succession we can represent M . We add a parity fix gadget to each box at the beginning of each wire, which will be useful later to ensure that each edge has been subdivided an even number of times. As a result, we have realized M using $O(1)$ bricks and $O(L) \times O(L) \times O(L)$ space. This collection of boxes is called a *general matching gadget*. A general matching gadget has a first and a last brick where it connects to the rest of the construction, we call these bricks endbricks.

36:16 How Does Object Fatness Impact Packing?



■ **Figure 6** A module with general matching gadgets of the interface and the core, with the simplified image of a brick-tree (in red).

If the goal is to realize a matching within a cell with vertex set V_x , then we can just create two copies of V_x (denoted by V'_x and V''_x), with a complete bipartite graph between them. For a matching edge $v_i v_j \in \binom{V_x}{2}$, we identify it with the edge $v'_i v''_j$. Then we realize the matching of this biclique using a general matching gadget.

The branching gadget

The *branching gadget* creates for all indices in $[L/8]^2$ a disjoint copy of a star on 4 vertices (that is, a vertex of degree 3 with its neighborhood of 3 isolated vertices). This gadget contains four bricks, and realizes $(L/8)^2$ disjoint stars. We use the first two bricks ($B^{(1)}$ and $B^{(2)}$) of the parallel matching gadget. The third brick B' is a translate of the first brick $B^{(1)}$ with the vector $(3, 2, L-1)$, i.e., the coordinates of $B'(i, j)$ are $(3i+3, 3j+2+i/L, L-1-3i)$. The final brick B'' is the translate of $B^{(2)}$ by the vector $(L, 0, 0)$. See Figure 5 for a rendering of the first “column” of the four bricks. Vertices corresponding to $B^{(2)}$ have degree three, and their neighbors are the boxes of the same index in $B^{(1)}$, B' and B'' .

Constructing a module

Our goal is to define modules of side length $O(L)$ that are capable of representing the role played by cells. The modules together must be able to represent a subgraph of \mathcal{C} of maximum degree three, where the neighbors of any vertex lie in distinct cells.

For all pairs of neighboring modules, we introduce a general matching gadget to represent the matching required by G between the two neighboring cells. These gadgets form the *interface*. Moreover, in the middle of each module, we add another general matching gadget to represent the matching within the cell; this gadget is the *core* of the module. See Figure 6. Finally, within each module, we tie the endbricks of the core and the endbricks of the interface falling inside the module together with a *brick-tree*. The brick-tree is a collection of $(L/8)^2$ isomorphic and disjoint trees, realized as a collection of branching, elbow, adjustment and bridge gadgets. Each tree (i, j) has maximum degree three, and its leaves are the boxes of index (i, j) in the interface and in the core.

First, we show that such a construction is sufficient to represent an even subdivision of an arbitrary subgraph G , and later we show how the brick-tree can be constructed. Let G be a subgraph with the desired properties, and let x be a particular cell. For each edge uv induced by x , we fix an arbitrary orientation, and realize the acquired matching so that the source vertex of the arcs are in one end of the core and the targets are in the other. Since the neighbors of any vertex lie in different cells, all indices of $[L/8]^2$ appear at most once, either as a source of an arc, as a target of an arc, or not at all. Then we realize the arcs using the core's general matching gadget of the module. For each index $\mathbf{i} \in [L/8]^2$, the edges incident to vertex \mathbf{i} of x can be assigned to a subtree T of the tree corresponding to index \mathbf{i} , where T has at most three leaves, at most one of which is adjacent to a box of the core, and other leaves are adjacent to boxes in distinct endbricks of the interface. There is a unique minimal subtree T that induces the desired (at most three) leaves; we can map a vertex $v \in V(G)$ of degree three to the degree three vertex of T . If V has a smaller degree, then it can be mapped to an arbitrary non-leaf vertex of T .

To construct a brick-tree in \mathbb{R}^3 , consider first a Euclidean grid cube of size $O(1)$. We can use this small cube as a model of our module: in general, an edge of this cube represents a brick. We have some edges already occupied by the general matching gadgets corresponding to the interface and the core. By choosing a cube large enough, we can ensure that these vertices are distant in the ℓ_1 norm. It is easy to see that if the cube is large enough (we allow its size to depend only on d), then there is a subtree of the grid of maximum degree three, where the leaves are some distant prescribed vertices. Such a tree can be constructed for example by mimicking a Hamiltonian path of the inscribed octahedron of the module, and adding to it small "spikes" that go to the endbrick of the interfaces. At the end of the path, we extend it towards the center of the cube, where we add another branching for the two endbricks of the core. The branching points in the brick-tree are branching gadgets, the turns are elbow gadgets, and straight segments are bridges and adjustments.

Finalizing the construction in \mathbb{R}^3

By packing the modules in a side length $O(sL)$ Euclidean cube, and removing unused boxes from each module according to the given subgraph, we get our final construction for three dimensions. For each edge, we have it represented by a sequence of $O(1)$ boxes passing through a single general matching gadget. Using the parity fix gadget inside the general matching gadget, we can ensure that the path representing the edge has an odd number of internal vertices. Therefore, the final construction has $O(|V(G)|)$ boxes, and each edge of G is represented with a path of odd length, that is, the graph induced by the boxes is an even subdivision of G .

The construction in higher dimensions

It is surprisingly easy to adapt our three-dimensional construction to the d -dimensional case. This time, we need to realize a subgraph of $\mathcal{C} = \mathcal{BEC}^d(s, (L/8)^{d-1})$.

The basic brick in d dimensions contains $(L/8)^{d-1}$ boxes, indexed by $[L/8]^{d-1}$, where the lexicographically minimal corner of box \mathbf{i} is $(0, 3\mathbf{i})$. For normal bricks, we allow perturbations of the form $3k$ ($|k| \in [L/8]$) along the axis of the brick, and k/L ($|k| \in [L/8]$) in all other directions. The parity fix, adjustment, and elbow gadgets can be defined analogously. The parallel matching gadget is also straightforward: the task here is to represent a parallel matching, where each edge is of the form $(\mathbf{i}, \mathbf{i}') \in [L/8]^{d-1} \times [L/8]^{d-1}$, where \mathbf{i}, \mathbf{i}' differ only on

36:18 How Does Object Fatness Impact Packing?

the t -th coordinate for some fixed $t \in [d-1]$. As previously, we can extend this to $(L/8)^{d-2}$ permutations, where for each $\iota \in [L/8]^{d-2}$, we have a permutation π_ι over the “column” ι , i.e., over the set

$$\{(i_1, \dots, i_{d-1}) \mid i_t \in [L/8] \text{ and } (i_1, \dots, i_{t-1}, i_{t+1}, \dots, i_{d-1}) = \iota\}.$$

Such a permutation can be represented as described before: we replace the role played by the x_1 axis with x_t , the role of x_2 with $x_{t+1 \bmod (d-1)}$ and x_3 with x_d . Along all other axes, we introduce no perturbations to the boxes. The column gadget corresponding to column $\iota = (i_1, \dots, i_{t-1}, i_{t+1}, \dots, i_{d-1})$ can be covered by¹

$$\begin{aligned} & [3i_1, 3i_1 + 1] \times \dots \times [3i_{t-1}, 3i_{t-1} + 1] \\ & \quad \times [-L/2, 3L/2] \times (3i_{t+1} - 0.5, 3i_{t+1} + 2.5) \\ & \quad \times [3i_{t+2}, 3i_{t+2} + 1] \times \dots \times [3i_{d-1}, 3i_{d-1} + 1] \times [0, \frac{3}{2}L]. \end{aligned}$$

These sets are clearly disjoint for distinct values of ι .

A general matching M is regarded as a permutation of $[L/8]^{d-1}$, which can be written as the product of $2(d-1) - 1$ special permutations by Corollary 12 that correspond to parallel matchings; therefore, M is realizable as the succession of $2d - 3$ parallel matchings. As a result, we can realize M with $O(d) = O(1)$ bricks and $O(L) \times \dots \times O(L)$ space. As before, we add parity fix gadgets to each box of one of the endbricks.

To realize a brick-tree, we can again trace a Hamiltonian path of the graph given by the dimension 1 faces of the cross-polytope inside the module, and add spikes to it to reach the endbricks of the interface and extend it to the two endbricks of the core. Note that the cross-polytope does have a Hamiltonian path, we can use e.g.

$$(1, 0, \dots, 0); (0, 1, 0, \dots, 0) \dots (0, \dots, 0, 1); (-1, 0, \dots, 0); (0, -1, 0, \dots, 0) \dots (0, \dots, 0, -1).$$

The finalizing steps are again analogous to the 3-dimensional case. This concludes the proof of Theorem 16.

¹ The formula is only accurate for the case $t \leq d - 2$. If $t = d - 1$, the role of x_{t+1} and x_1 should be switched.