

On Optimization and Counting of Non-Broken Bases of Matroids

Dorna Abdolazimi ✉

University of Washington, Seattle, WA, USA

Kasper Lindberg ✉

University of Washington, Seattle, WA, USA

Shayan Oveis Gharan ✉

University of Washington, Seattle, WA, USA

Abstract

Given a matroid $M = (E, \mathcal{I})$, and a total ordering over the elements E , a broken circuit is a circuit where the smallest element is removed and an NBC independent set is an independent set in \mathcal{I} with no broken circuit. The set of NBC independent sets of any matroid M define a simplicial complex called the broken circuit complex which has been the subject of intense study in combinatorics. Recently, Adiprasito, Huh and Katz showed that the face of numbers of any broken circuit complex form a log-concave sequence, proving a long-standing conjecture of Rota.

We study counting and optimization problems on NBC bases of a generic matroid. We find several fundamental differences with the independent set complex: for example, we show that it is NP-hard to find the max-weight NBC base of a matroid or that the convex hull of NBC bases of a matroid has edges of arbitrary large length. We also give evidence that the natural down-up walk on the space of NBC bases of a matroid may not mix rapidly by showing that for some family of matroids it is NP-hard to count the number of NBC bases after certain conditionings.

2012 ACM Subject Classification Theory of computation → Random walks and Markov chains; Theory of computation → Computational complexity and cryptography

Keywords and phrases Complexity, Hardness, Optimization, Counting, Random walk, Local to Global, Matroids

Digital Object Identifier 10.4230/LIPIcs.APPROX/RANDOM.2023.40

Category RANDOM

Funding *Dorna Abdolazimi*: Research supported by NSF grant CCF-2203541, and Air Force Office of Scientific Research grant FA9550-20-1-0212.

Kasper Lindberg: Research supported by NSF grant CCF-2203541.

Shayan Oveis Gharan: Research supported by NSF grant CCF-2203541, Air Force Office of Scientific Research grant FA9550-20-1-0212, and Simons Investigator grant.

1 Introduction

A matroid $M = (E, \mathcal{I})$ consists of a finite ground set E and a collection \mathcal{I} of subsets of E , called independent sets, satisfying:

Downward closure: If $S \subseteq T$ and $T \in \mathcal{I}$, then $S \in \mathcal{I}$.

Exchange axiom: If $S, T \in \mathcal{I}$ and $|T| > |S|$, then there exists an element $i \in T \setminus S$ such that $S \cup \{i\} \in \mathcal{I}$.

The rank of a set $S \subseteq E$ is the size of the largest independent set contained in S . All maximal independent sets of M , called the bases of M , have the same size r , which is called the rank of M .



© Dorna Abdolazimi, Kasper Lindberg, and Shayan Oveis Gharan;
licensed under Creative Commons License CC-BY 4.0

Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2023).

Editors: Nicole Megow and Adam D. Smith; Article No. 40; pp. 40:1–40:14



Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

Sampling and counting problems on matroids have captured the interest of many researchers for several decades with applications to reliability [6], liquidity of markets [19], etc. A recent breakthrough in this field proved that the down-up walk on the bases of a matroid mixes rapidly to the (uniform) stationary distribution and can be used to count the number of bases of a matroid [3, 7], resolving the conjecture of Mihail and Vazirani from 1989 [18]. The down-up walk is easy to describe: Start with an arbitrary base B and repeatedly execute the following two steps:

1. Choose a uniformly random element $i \in B$ and delete it.
2. Among all bases (of M) that contain $B \setminus \{i\}$, choose one uniformly at random.

A central question that has puzzled researchers since then is sampling a non-broken (circuit) basis (NBC basis) of a matroid [4].

A set $C \subseteq E$ is a *circuit* iff $C \setminus \{e\} \in \mathcal{I}$ for any $e \in C$. A *broken circuit* (with respect to a total ordering \mathcal{O}) is a set $C \setminus \{e\}$, where $C \subseteq E$ is a circuit and e is the **smallest** element of C with respect to \mathcal{O} . An independent set $S \subseteq E$ is a *non-broken* independent set (NBC independent set) if it contains no broken circuits. The NBC independent sets are closely related to several interesting combinatorial objects. The number of NBC independent sets of size k in a graphic matroid is equal to the absolute value of the $(n - 1) - k$ -th coefficient of the chromatic polynomial of the underlying graph where n is the number of vertices. As a corollary the following facts hold:

► **Fact 1.** *The following facts are well-known about the counts of NBC bases/independent sets of different family of matroids.*

- *The number of all NBC independent sets of a graphic matroid is equal to the the number of acyclic orientations of the graph [22].*
- *The number of all NBC independent sets of a co-graphic matroid is equal to the number of strongly connected orientations of the graph (see e.g., [13]).*
- *The number of non-broken spanning trees of a graph is equal to the number of parking functions with respect to a unique source vertex [4]*
- *The number of NBC independent of sets of linear matroid with vectors v_1, \dots, v_n is equal to the number of regions defined by the intersection of the orthogonal hyperplanes (see e.g., [23]).*

We emphasize that although the set of NBC independent sets/bases of a matroid are functions of the underlying total order \mathcal{O} , the counts of the number NBC independent sets of rank k for any $0 \leq k \leq r$ are invariant under \mathcal{O} [23]. We remark that, to the best of our knowledge as of this date, none of the above counting problems are known to be computationally tractable.

Given a matroid M with an arbitrary total ordering \mathcal{O} , one can analogously run the down-up walk only on the NBC bases of M . It is not hard to see that this chain is irreducible and converges to the uniform stationary distribution. Following the work of [3] it was conjectured that the down-up walk on the NBC bases of any matroid mixes rapidly ¹.

► **Conjecture 2.** *For any matroid M , and any total ordering \mathcal{O} of the elements of M , the down-up walk on the NBC bases of a matroid mixes in polynomial time.*

¹ In fact, this conjecture was raised as an open problem in several recent workshops UC Santa Barbara workshop on New tools for Optimal Mixing of Markov Chains: Spectral Independence and Entropy Decay, and Simon's workshop on Geometry of Polynomials

It turns out that the above conjecture, if true, would be give a generalization of the result of [3], because of the following fact.

► **Fact 3** ([5]). *For any matroid M one can construct another matroid M' with an ordering \mathcal{O} with only one extra element such that there is a bijection between bases of M and non-broken bases of M' .*

Furthermore, if the above conjecture is true, then since matroids are closed under truncation, one can also count the number of all NBC independent sets of M , thus resolving all of the open problems in Fact 1.

A promising reason to expect these problems to be tractable in the first place is the remarkable work of Adiprasito, Huh and Katz [1] who proved the Rota's conjecture showing that the face numbers of a broken circuit complex (see below for definition) of any matroid forms a log-concave sequence. For comparison, it is well-known that the coefficients of the matching polynomial of any graph form a log-concave sequence and the classical algorithm of Jerrum-Sinclair [15] gives an efficient algorithm to count the number of matchings of any graph (although to this date we still don't know an efficient algorithm to count the number of **perfect** matchings of general graphs).

1.1 Background

The existing analyses of the mixing time of the down-up walk for bases of matroids, crucially rely on the theory of high dimensional simplicial complexes [3, 17], which has found many intriguing applications in several areas of computer science and math in the past few years [14].

Simplicial Complex. A *simplicial complex* X on a finite ground set U is a downwards closed set system, i.e. if $\tau \in X$ and $\sigma \subset \tau \subseteq U$, then $\sigma \in X$. The elements of X are called faces, and the maximal faces are called facets. We say X is a pure d -dimensional complex if all of its facets are of size d . We denote the set of facets by $X(d)$. A weighted simplicial complex (X, π) is a simplicial complex X paired with a probability distribution π on its facets. The global walk (down-up walk) P^\vee on the facets of a d -dimensional complex (X, π) is defined as follows: starting at a facet τ , we transition to the next facet τ' by the following two steps:

1. Select a uniformly random element $x \in \tau$ and remove x from τ .
2. Select a random facet τ' containing $\tau \setminus \{x\}$ with probability proportional to $\pi(\tau')$.

Broken Circuit Complex. For a concrete example, it turns out that the set NBC independent sets of any matroid M (with respect to any ordering \mathcal{O}) form a **pure** simplicial complex that is known as the *broken circuit complex*. We denote this complex by $BC(M, \mathcal{O})$. We state purity as the following fact.

► **Fact 4.** *For every NBC independent set I , there exists an NBC base B such that $I \subseteq B$.*

The face numbers of the complex $BC(M, \mathcal{O})$ is the sequence n_0, n_1, \dots, n_r where n_i is the number of NBC independent sets of rank i . As alluded to above this sequence is in variant over \mathcal{O} . The down-up walk over this complex equipped with a uniform distribution over its facets is the same as the down-up walk over NBC bases we explained before.

The link of a face $\tau \in X$ is the simplicial complex $X_\tau := \{\sigma \setminus \tau : \sigma \in X, \sigma \supset \tau\}$. For each face τ , we define the induced distribution π_τ on the facets of X_τ as

$$\pi_\tau(\eta) = \Pr_{\sigma \sim \pi} [\sigma \supset \eta \mid \sigma \supset \tau]. \quad (1)$$

Local Walks. For any face τ of size $0 \leq k \leq d - 2$, the local walk for τ is a Markov chain on the ground set of X_τ with transition probability matrix P_τ is defined as

$$P_\tau(x, y) = \frac{1}{d - k - 1} \Pr_{\sigma \sim \pi_\tau} [y \in \sigma \mid \sigma \supset \tau \cup \{x\}]. \quad (2)$$

for distinct x, y in the ground set of X_τ .

The following theorem shows that the spectral expansion of the global walk P^\vee on a simplicial complex can be bounded through bounding the local spectral expansion of the complex.

► **Theorem 5** (Local-to-Global Theorem [10, 17, 9, 2]). *Say a d -dimensional weighted simplicial complex (X, π) is a $(\gamma_0, \dots, \gamma_{d-2})$ -local spectral expander if for every face τ of size $0 \leq k \leq d - 2$, the second largest eigenvalue of P_τ is at most γ_k , i.e., $\lambda_2(P_\tau) \leq \gamma_k$.*

Given a weighted simplicial complex (X, π_d) that is a $(\gamma_0, \dots, \gamma_{d-2})$ -local spectral expander, the down-up walk which samples from π has spectral gap lower bounded by

$$1 - \lambda_2(P^\vee) \geq \frac{1}{d} \prod_{j=0}^{d-2} (1 - \gamma_j)$$

To prove that the down-up walk mixes rapidly on the bases of any matroid, [3] proved that the independent set complex of any matroid M is a $(0, 0, \dots, 0)$ -local spectral expander. Building on this, a natural method to prove Conjecture 2 is to show that the broken circuit complex of any matroid M of rank r and for any total ordering is a $(\gamma_0, \dots, \gamma_{r-2})$ -local spectral expander for $\gamma_i \leq \frac{O(1)}{r-i}$.

► **Conjecture 6.** *For any matroid M of rank r and any ordering \mathcal{O} the broken circuit complex of M is a $(\gamma_0, \dots, \gamma_{r-2})$ -local spectral expander for some $\gamma_i \leq \frac{O(1)}{r-i}$*

1.2 Our results

Our main result is to disprove Conjecture 6 in a very strong form, namely for the class of (truncated) graphic matroids.

► **Theorem 7.** *There exists an infinite sequence of (truncated) graphic matroids M_1, M_2, \dots with orderings $\mathcal{O}_1, \mathcal{O}_2, \dots$, such that for every $n \geq 1$, M_n has $\text{poly}(n)$ elements, and there exists a face τ of the broken circuit complex of $X = \text{BC}(M_n, \mathcal{O})$ for which the down-up walk on the facets of the link X_τ has a spectral gap of at most $n^{-\Omega(n)}$.*

In fact, we even prove a stronger statement

► **Theorem 8.** *Given a matroid $M = (E, \mathcal{I})$ and a total ordering \mathcal{O} and a set $S \subseteq E$, unless $\text{RP} = \text{NP}$, there is no FPRAS for counting the number of NBC bases of M that contain S .*

Although this theorem does not refute Conjecture 2, it shows that one probably need different techniques (or probably a different chain) to sample/count NBC bases of a matroid. Indeed, one may even need a different proof for the performance of down-up walk to sample ordinary bases of matroids.

To complement our main results we also prove that, unlike optimization on bases of a matroid, optimization is NP-hard on the NBC bases of matroids. Moreover, unless $\text{NP} = \text{RP}$, there is no FPRAS for computing the sum of the weights of all NBC bases of a matroid subject to an external field, while the same computation over the bases of a matroid has a FPRAS.

► **Theorem 9.** *Given a matroid $M = (E, \mathcal{I})$ with $|E| = n$ elements, an arbitrary total ordering \mathcal{O} , and weights w_1, \dots, w_n , it is NP-hard to find the maximum weight NBC basis of M , where the weight of a NBC basis B is $\sum_{i \in B} w_i$.*

► **Theorem 10.** *Given a matroid $M = (E, \mathcal{I})$ with $|E| = n$ elements, a total ordering \mathcal{O} , and weights $\{1 \leq \lambda_e \leq O(n)\}_{e \in E}$, unless $NP = RP$, there is no FPRAS for computing the partition function of the λ -external field applied to uniform distribution of NBC independent sets, i.e., there is no FPRAS for computing :*

$$\sum_{B \text{ NBC Base}} \prod_{e \in B} \lambda_e.$$

It is well known that a 0/1-polytope (i.e. the convex hull of a subset $S \subseteq \{0, 1\}^n$) has all vertices of equal hamming weight r and edges of ℓ_2 length $\sqrt{2}$ iff the polytope is a matroid base polytope of rank r [12]. Moreover, assuming the Mihail-Vazirani conjecture, there is efficient algorithm to sample a uniformly random vertex of a 0/1-polytope with constant sized edge length [18].

We show that, unlike matroids, the NBC Base polytope, i.e. the convex hull of the indicator vectors of all NBC bases of a matroid M , has edges of arbitrarily long length.

► **Theorem 11.** *For any n , there exists a graphic matroid M with n elements and a total ordering \mathcal{O} such that the convex hull of all NBC bases of M has edges of ℓ_2 length at least $\Omega(\sqrt{n})$.*

2 Preliminaries

Given a graph $G = (V, E)$, we denote the number of independent sets of size i of G by $i_k(G)$. For every set $S \subseteq V$, we define $N(S) := \{v \notin S : \exists u \in S, \{u, v\} \in E\}$ as the set of neighbors of S in G .

► **Definition 12 (Conductance).** *Given a weighted d -regular graph $G = (V, E, w)$, with weights $w : E \rightarrow \mathbb{R}_{\geq 0}$, for $S \subseteq V$, the conductance of S is defined as*

$$\phi(S) = \frac{w(S, \bar{S})}{d|S|},$$

where $w(S, \bar{S})$ is the sum of the weights of edges in the cut (S, \bar{S}) . Note that since G is regular, the weighted degree of every vertex is d . The conductance of G is defined as

$$\phi(G) = \min_{S: |S| \leq |V|/2} \phi(S).$$

Given a weighted graph $G = (V, E, w)$, the simple random walk is the following stochastic process: Given $X_0 = v \in V$, for every $u \sim v$, we have $X_1 = u$ with probability $\frac{w_{\{u,v\}}}{d_w(v)}$ and we let P be the transition probability matrix of the walk.

The following theorem is well-known and follows from the easy side of the Cheeger's inequality.

► **Theorem 13.** *For any regular graph $G = (V, E)$ and any set $S \subseteq V$ and $|S| \leq |V|/2$*

$$\frac{1 - \lambda_2(P)}{2} \leq \phi(G) \leq \phi(S) \leq \frac{|N(S)|}{|S|}$$

where $1 - \lambda_2(P)$ is the spectral gap of the simple random walk on G .

40:6 On Optimization and Counting of Non-Broken Bases of Matroids

A graphic matroid $M = (E, \mathcal{I})$ is a matroid defined on the edges of a graph $G = (V, E)$ and its independent sets are all subsets of edges that do not contain any cycle. It is easy to verify that circuits of M correspond to cycles of G .

► **Definition 14** (Matroid Truncation). *Let $M = (E, \mathcal{I})$ be a matroid of rank r . The truncation of M to rank $r' \leq r$ removes all independent sets of size strictly greater than r' . It is easy to see that the truncation of any matroid M to any $r' \leq r$ is also a matroid.*

Let M' be the truncation to rank r' of a graphic matroid of rank r defined on the edges of a graph G . The bases of M' correspond to forests with r' edges and the circuits of M' are the circuits of G along with all spanning forests of size $r' + 1$.

The following fact about polytopes follows from convexity.

► **Fact 15.** *For any polytope $P \subseteq \mathbb{R}^d$ with vertices $v_1, \dots, v_n \in \mathbb{R}^d$, $\{v_i, v_j\}$ is an edge of P iff there exists a weight function $w \in \mathbb{R}^d$ such that*

$$\langle w, v_i \rangle = \langle w, v_j \rangle > \langle w, v_k \rangle,$$


for any $k \neq i, j$.

3 Results

We start with proving Theorem 11.

► **Theorem 11.** *For any n , there exists a graphic matroid M with n elements and a total ordering \mathcal{O} such that the convex hull of all NBC bases of M has edges of ℓ_2 length at least $\Omega(\sqrt{n})$.*

Proof. Let n be odd. Consider the following graphic matroid M (with n edges), with the ordering \mathcal{O} : $1 < 2 < \dots < n$ defined by the edges of the following graph:



We show that for $B = \{n\} \cup \{2i - 1 : 1 \leq i \leq \frac{n-1}{2}\}$ and $B' = \{1\} \cup \{2i : 1 \leq i \leq \frac{n-1}{2}\}$, $\{B, B'\}$ forms an edge in the NBC matroid base polytope denoted as P_M . We define a $w \in \mathbb{R}^n$ and then use Fact 15 to prove the statement. Let $w_n = \frac{n+1}{2}$, and for any $1 \leq i \leq \frac{n-1}{2}$, let $w_{2i} = 1$ and $w_{2i-1} = 0$. It is easy to check that the function $\langle w, \mathbf{1}_B \rangle = \langle w, \mathbf{1}_{B'} \rangle = \frac{n+1}{2}$ and $\langle w, \mathbf{1}_{B''} \rangle < \frac{n+1}{2}$ for all NBC basis $B'' \neq B, B'$. Therefore $\{B, B'\}$ forms an edge in P_M . The statement follows from the fact that $\|\mathbf{1}_B - \mathbf{1}_{B'}\|_2 = \sqrt{n}$. ◀

Next, we prove Theorem 9 via a reduction from the MAX-INDEP-SET problem: Given a graph $G = (V, E)$, a weight function $w : V \rightarrow \mathbb{R}_{\geq 0}$, and an integer k , decide whether G has an independent set of weight at least k or not.

Note that independent sets of G and independent sets of a BC complex/matroid are two different notions. To complete the proof we use the following well-known hardness result.

► **Theorem 16** ([16]). *MAX-INDEP-SET is NP-complete.*

► **Theorem 9.** *Given a matroid $M = (E, \mathcal{I})$ with $|E| = n$ elements, an arbitrary total ordering \mathcal{O} , and weights w_1, \dots, w_n , it is NP-hard to find the maximum weight NBC basis of M , where the weight of a NBC basis B is $\sum_{i \in B} w_i$.*

Proof. We prove this by a reduction from MAX-INDEP-SET. Let $G = (V, E)$ be a graph, a vertex weight function $w : V \rightarrow \mathbb{R}_{\geq 0}$ and k an integer. Construct a new graph $G' = (V', E')$ from G by first copying G and then adding a new vertex z and edges $e_v = \{z, v\}$ for all $v \in V$. We define $w' : E' \rightarrow \mathbb{R}_{\geq 0}$ as $w'(e_v) = w(v)$ for every $v \in V$, and $w'(e) = 0$ for every $e \in E$. Moreover, consider the following total ordering \mathcal{O} on E' :

$$E < \{e_v : v \in V\},$$

where the ordering within each set is arbitrary. Let M be the graphic matroid defined by the edges of G' , we will be look at bases/independent sets of $\text{BC}(M, \mathcal{O})$.

▷ **Claim 17.** There exists an independent set of G of weight at least k iff there exists an NBC basis of M with weight at least k .

We prove the claim in a straightforward manner. Suppose there is an independent set $I \subseteq V$ of G with $w(I) \geq k$ and consider the set $I' \subseteq E'$ defined by $I' = \{e_v : v \in I\}$. By definition, $w'(I') \geq k$. We argue that I' does not contain any broken circuit. Assume otherwise that there is a broken circuit $C \setminus \{e\} \subseteq I'$. Since C corresponds to a cycle in G' and $C \setminus \{e\}$ is contained in I' , it is not hard to see that $C \setminus \{e\} = \{e_v, e_{v'}\}$ for some $v, v' \in I$ and $e = \{v, v'\}$ is an edge in G . But this is a contradiction with the fact that I is an independent set of G .

Hence I' is a NBC independent set. Since the broken circuit complex is pure (see Fact 4), there exists an NBC basis B containing I' which has weight $w'(B) \geq w'(I') \geq k$.

For the other direction, suppose we have a NBC basis $B' \subseteq E'$ of weight $w'(B') \geq k$, and define $I \subseteq V$ by $I = \{v : e_v \in B'\}$. Since all edges coming from E have zero weight, $w(I) = w'(B') \geq k$. To see that I is an independent set of G' , note that if there is an edge $\{v, v'\}$ for some $v, v' \in I$, we have $e_v, e_{v'} \in B'$, then $\{e_v, e_{v'}\}$ forms a broken circuit according to the ordering \mathcal{O} . Therefore I is an independent set of G of weight at least k . ◀


It's important to note that the above proof works under the crucial assumption that the order \mathcal{O} is chosen carefully based on the weights (and in some sense in the same order of the weights).

We can amplify the ideas in the previous construction to also argue Theorem 7. This is done by constructing a Broken Circuit complex for which the down-up walk of a carefully chosen link has inverse exponentially small spectral gap.

► **Theorem 7.** *There exists an infinite sequence of (truncated) graphic matroids M_1, M_2, \dots with orderings $\mathcal{O}_1, \mathcal{O}_2, \dots$, such that for every $n \geq 1$, M_n has $\text{poly}(n)$ elements, and there exists a face τ of the broken circuit complex of $X = \text{BC}(M_n, \mathcal{O})$ for which the down-up walk on the facets of the link X_τ has a spectral gap of at most $n^{-\Omega(n)}$.*

Proof. Take the complete bipartite graph $G = K_{n,n} = (A, B, E = A \times B)$, with $|A| = |B| = n$. Also, let $V = A \cup B$. Let $\ell \geq 1$ be a parameter that we choose later, and construct a new graph

$$G' = (V' = V \cup \{y, z\} \cup \{z_{v,i} : v \in V, i \in [\ell]\}, E' = E \cup \{e_0\} \cup \{e_{v,i}, f_{v,i} : v \in V, i \in [\ell]\})$$



■ **Figure 1** A schematic of the graph G' in the proofs of Theorem 7 and Theorem 8 where $G = K_{n,n}$ is the complete bipartite graph in the former and it is a hard instance of $\sharp\text{INDEP-SET-INC}(7, \frac{2}{19})$ in latter.

where $e_0 = \{y, z\}$, $e_{v,i} = \{z, z_{v,i}\}$, $f_{v,i} = \{z_{v,i}, v\}$ (see Figure 1). For a sanity check, note that $|V| = 2n$ and $|V'| = 2\ell n + 2n + 2$.

Let $M = (E', \mathcal{I})$ be the graphic matroid defined by G' truncated to rank $2\ell n + n + 1$, i.e., the bases of M are forests of G' with exactly $2\ell n + n + 1$ edges. Now, consider the following total ordering \mathcal{O} on E' :

$$e_0 < E < \{e_{v,i} : v \in V, i \in [\ell]\} < \{f_{v,i} : v \in V, i \in [\ell]\},$$

where the ordering within each set is arbitrary.

Moreover, let $X := \text{BC}(M, \mathcal{O})$, and define

$$\tau = \{e_{v,i} : v \in V, i \in [\ell]\}.$$

For simplicity of notation, let $F_A := \{f_{v,i} : v \in A, i \in [\ell]\}$ and $F_B := \{f_{v,i} : v \in B, i \in [\ell]\}$.

▷ **Claim 18.** For any facet S of X_τ , either $S \cap F_A = \emptyset$, or $S \cap F_B = \emptyset$,

This follows from the fact that G is a complete bipartite graph and edges in E are smaller than $e_{v,i}$'s and $f_{u,j}$'s; so if $S \cap F_A, S \cap F_B \neq \emptyset$, then it has a broken circuit.

Therefore, the set of facets of X_τ can be partitioned into $2n + 1$ sets $(\cup_{i=1}^n \mathcal{S}_{A,i}) \cup (\cup_{i=1}^n \mathcal{S}_{B,i}) \cup \mathcal{S}_0$, where $\mathcal{S}_{A,i}$ is the set of all facets S with $|S \cap F_A| = i$, $\mathcal{S}_{B,i}$ is the set of all facets S with $|S \cap F_B| = i$, and \mathcal{S}_0 is the set of all facets with $|S \cap (F_A \cup F_B)| = 0$. Let $\mathcal{S}_A := \cup_{i=1}^n \mathcal{S}_{A,i}$ and similarly define \mathcal{S}_B . We show that $\frac{|N(\mathcal{S}_A)|}{|\mathcal{S}_A|} \leq n^{-\Omega(n)}$, where $N(\mathcal{S}_A)$ is the set of neighbors of \mathcal{S}_A in the down-up walk P_τ^\vee on the facets of τ . WLOG we can assume that $|\mathcal{S}_A|$ is at most half of all facets. Applying Theorem 13, this would imply that $1 - \lambda_2(P_\tau^\vee) \leq n^{-\Omega(n)}$.

First, note that for every facet $S \in \mathcal{S}_A$ and $T \in \mathcal{S}_B \setminus \mathcal{S}_{B,1}$, we get $P^\vee(S, T) = 0$ since $|S \Delta T| > 2$. So, $N(\mathcal{S}_A) \subseteq \mathcal{S}_{B,1} \cup \mathcal{S}_0$. First, notice $|\mathcal{S}_0| \leq \binom{|E|}{n} \leq n^{2n}$. Furthermore, $|\mathcal{S}_{B,1}| \leq \binom{n}{1} \ell \binom{|E|}{n-1} \leq \ell n^{2n}$.

This follows from the fact that any facet in $\mathcal{S}_{B,1}$ can be written as $\{f_{v,i_v}\} \cup \{e_0\} \cup K$ for some $v \in A$, $i_v \in [\ell]$, and subset $K \subseteq E$ of size $n - 1$.

Lastly, $|\mathcal{S}_A| \geq |\mathcal{S}_{A,n}| = \ell^n$. This is because every choice of $\{i_v\}_{v \in A}$ corresponds to a set in $\mathcal{S}_{A,n}$ whose sets are of the form $\{f_{v,i_v} : v \in V\} \cup \{e_0\}$. These sets all don't contain a broken circuit because the circuits introduced through truncation are exactly the forests with

$2\ell n + n + 2$ edges. However, any proper superset of $\{f_{v,i_v} : v \in V\} \cup \{e_0\}$ must include e_0 , so looking at the circuit introduced by the superset, the corresponding broken circuit will always remove e_0 . Putting it all together,

$$1 - \lambda_2(P_\tau^\vee) \leq \frac{|N(\mathcal{S}_A)|}{|\mathcal{S}_A|} \leq \frac{n^{2n}(1+\ell)}{\ell^n} \stackrel{\text{assuming } \ell \geq n^3}{\leq} n^{-\Omega(n)}$$

as desired. \blacktriangleleft

We prove Theorem 10 and Theorem 8 by a reduction from $\sharp\text{INDEP-SET-INC}(7, \frac{2}{19})$, defined as the following.

► Definition 19 ($\sharp\text{INDEP-SET-INC}(7, \frac{2}{19})$). *Given a 7-regular graph $G = (V, E)$ that satisfies $i_k(G) \leq i_{\lfloor \frac{2|V|}{19} \rfloor}(G)$ for any $k < \lfloor \frac{2|V|}{19} \rfloor$, where $i_k(G)$ are the independent sets of G of size k , count the number of independent sets of size $\lfloor \frac{2|V|}{19} \rfloor$.*

► Theorem 20. *Unless $\text{NP} = \text{RP}$, there is no randomized algorithm with constant approximation ratio for $\sharp\text{INDEP-SET-INC}(7, \frac{2}{19})$.*

We leave the proof of this for the appendix.

Now, we are ready to prove Theorem 8. The high-level structure of the proof is similar to the proof of Theorem 7 where we apply a similar gadget to graphs on which it is hard to count independent sets (as opposed to the complete bipartite graph).

► Theorem 8. *Given a matroid $M = (E, \mathcal{I})$ and a total ordering \mathcal{O} and a set $S \subseteq E$, unless $\text{RP} = \text{NP}$, there is no FPRAS for counting the number of NBC bases of M that contain S .*

Proof. For simplicity of notion, let $\alpha := \frac{2}{19}$. We prove by a reduction from $\sharp\text{INDEP-SET-INC}(7, \frac{2}{19})$. Take any arbitrary 7-regular graph $G = (V, E)$ whose number of independent sets of size $\lfloor \alpha|V| \rfloor$ is at least the number of its independent sets of size k for any $k < \lfloor \alpha|V| \rfloor$. Let $n := |V|$ and N be the number of independent sets of size $\lfloor \alpha n \rfloor$ of G . Also, define $\ell \geq 1$ to be a parameter that we choose later.

Now, construct a new graph

$$G' = (V' = V \cup \{y, z\} \cup \{z_{v,i} : v \in V, i \in [\ell]\}, E' = E \cup \{e_0\} \cup \{e_{v,i}, f_{v,i} : v \in V, i \in [\ell]\})$$

where $e_0 = \{y, z\}$, $e_{v,i} = \{z, z_{v,i}\}$, $f_{v,i} = \{z_{v,i}, v\}$ (see Figure 1). Let $M = (E', \mathcal{I})$ be the graphic matroid defined by G' truncated at rank $\ell n + \lfloor \alpha n \rfloor + 1$, i.e., the bases of M are forests of G' with exactly $\ell n + \lfloor \alpha n \rfloor + 1$ edges. Now, consider the following ordering \mathcal{O} on E' :

$$e_0 < E < \{e_{v,i} : v \in V, i \in [\ell]\} < \{f_{v,i} : v \in V, i \in [\ell]\},$$

where the ordering within each set is arbitrary. Moreover, let $X := \text{BC}(M, \mathcal{O})$, and define

$$\tau = \{e_{v,i} : v \in V, i \in [\ell]\}.$$

We claim that the number of facets of X_τ is at least $\ell^{\lfloor \alpha n \rfloor} N$ and at most $2\ell^{\lfloor \alpha n \rfloor} N$. So, a 1.5-approximation to the number facets of X_τ , i.e., the number NBC bases of M that contain τ , gives a 3-approximation to N , the number of independent sets of size $\lfloor \alpha n \rfloor$ of G .

We use the following crucial observation:

► Claim 21. For any facet S of X_τ , $\{v : \exists f_{v,i} \in S\}$ is an independent set of G and for any $f_{v,i}, f_{v,j} \in S$ we have $i = j$.

Conversely, for any $S \subseteq \{f_{v,i} : v \in V, i \in [\ell]\}$, such that the set $\{v : \exists f_{v,i} \in S\}$ is an independent set of size $\lfloor \alpha n \rfloor$ of G , and $f_{v,i}, f_{v,j} \in S \implies i = j$, we have $S \cup \{e_0\}$ is a facet of X_τ .

40:10 On Optimization and Counting of Non-Broken Bases of Matroids

The proof simply follows from the fact that edges of E are smaller than $e_{v,i}$'s, and $f'_{u,j}$'s in \mathcal{O} . By the second part of the claim, we can write

$$|X_\tau(\lfloor \alpha n \rfloor + 1)| = \ell^{\lfloor \alpha n \rfloor} N + |\{S \in X_\tau(\lfloor \alpha n \rfloor + 1) : S \cap E \neq \emptyset\}| \geq \ell^{\lfloor \alpha n \rfloor} N. \quad (3)$$

Define $i_k := i_k(G)$ as the number of independent sets of size k of graph G . By the first part of the above claim we can write,

$$|\{S \in X_\tau(\lfloor \alpha n \rfloor + 1) : S \cap E \neq \emptyset\}| \leq \sum_{k=0}^{\lfloor \alpha n \rfloor - 1} \ell^k \cdot i_k \cdot \binom{|E|}{\lfloor \alpha n \rfloor - k} \quad (4)$$

$$\leq \sum_{k=0}^{\lfloor \alpha n \rfloor - 1} \ell^k \cdot i_k \cdot |E|^{\lfloor \alpha n \rfloor - k} \quad (5)$$

$$\stackrel{\text{using } i_k \leq N}{\leq} N |E|^{\lfloor \alpha n \rfloor} \sum_{k=0}^{\lfloor \alpha n \rfloor - 1} (\ell/|E|)^k \quad (6)$$

$$\stackrel{\text{assuming } \ell \geq 2|E|}{\leq} N |E|^{\lfloor \alpha n \rfloor} (\ell/|E|)^{\lfloor \alpha n \rfloor} \leq N \ell^{\lfloor \alpha n \rfloor} \quad (7)$$

Putting these together with (3) concludes the proof. \blacktriangleleft

► Theorem 10. *Given a matroid $M = (E, \mathcal{I})$ with $|E| = n$ elements, a total ordering \mathcal{O} , and weights $\{1 \leq \lambda_e \leq O(n)\}_{e \in E}$, unless $NP = RP$, there is no FPRAS for computing the partition function of the λ -external field applied to uniform distribution of NBC independent sets, i.e., there is no FPRAS for computing :*

$$\sum_{B \text{ NBC Base}} \prod_{e \in B} \lambda_e.$$

Proof. For simplicity of notion, let $\alpha := \frac{2}{19}$. The proof is similar to the proof of Theorem 8 by a reduction from $\sharp\text{INDEP-SET-INC}(7, \frac{2}{19})$. Take any arbitrary 7-regular graph $G = (V, E)$ with $n := |V|$ vertices whose number of independent sets of size $\lfloor \alpha |V| \rfloor$ is at least the number of its independent sets of size k for any $k < \lfloor \alpha |V| \rfloor$. Construct a new graph

$$G' = (V' = V \cup \{y, z\}, E' = E \cup \{e_0 = \{y, z\}\} \cup \{e_v = \{v, z\} : v \in V\})$$

Let $M = (E', \mathcal{I})$ be the graphic matroid given by G' truncated to rank $\lfloor \alpha n \rfloor + 1$ and consider the following ordering \mathcal{O} on E'' :

$$e_0 < E < \{e_v : v \in V\},$$

where as usual the ordering within each set is arbitrary. Define weights $\lambda : E' \rightarrow \mathbb{R}_{\geq 0}$ as follows:

$$\lambda_e = \begin{cases} \ell & \text{if } e = e_v \text{ for some } v \in V, \\ 1 & \text{o.w.} \end{cases},$$

for some ℓ that we choose later. We argue that

$$\lambda^{\lfloor \alpha n \rfloor} N \leq \sum_B \prod_{e \in B} \lambda_e \leq 2 \lambda^{\lfloor \alpha n \rfloor} N.$$

where here (and henceforth) the sum is over B 's that are NBC bases of M , and therefore a 1.5-approximation to the partition function, i.e., the quantity in the middle, is a 3-approximation to N . Similar to the previous theorem we have the following claim.

▷ Claim 22. For any NBC base B of M , we have $\{v : e_v \in B\}$ is an independent set of G . Conversely, for any independent set I of G of size $|I| = \lfloor \alpha n \rfloor$, $\{e_0\} \cup \{e_v : v \in I\}$ is a NBC base of M .

So,

$$\begin{aligned} \sum_B \prod_{e \in B} \lambda_e &= \sum_{B: B \cap E \neq \emptyset} \prod_{e \in B} \lambda_e + \sum_{B: B \cap E = \emptyset} \prod_{e \in B} \lambda_e \\ &= \sum_{B: B \cap E \neq \emptyset} \prod_{e \in B} \lambda_e + \ell^{\lfloor \alpha n \rfloor} |\{S \subseteq V : S \text{ independent set of } G, |S| = \lfloor \alpha n \rfloor\}| \end{aligned} \quad (8)$$

Define i_k as the number of independent sets of size k of graph G . We have

$$\sum_{B: B \cap E \neq \emptyset} \prod_{e \in B} \lambda_e \leq \sum_{k=0}^{\lfloor \alpha n \rfloor - 1} \ell^k i_k \binom{|E|}{\lfloor \alpha n \rfloor - k} \stackrel{\substack{\leq \\ \text{using } i_k \leq N, \\ \text{assuming } \ell \geq 2|E|}}{\leq} \ell^{\lfloor \alpha n \rfloor} N$$

where the last inequality follows from the same calculations as in Equation (4). ◀

References

- 1 Karim Adiprasito, June Huh, and Eric Katz. Hodge theory for combinatorial geometries, 2018. [arXiv:1511.02888](#).
- 2 Vedat Levi Alev and Lap Chi Lau. Improved analysis of higher order random walks and applications. In *Proceedings of the 52nd Annual ACM Symposium on Theory of Computing (STOC)*, 2020.
- 3 Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant. Log-concave polynomials ii: high-dimensional walks and an fpras for counting bases of a matroid. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 1–12, 2019.
- 4 Brian Benson, Deeparnab Chakrabarty, and Prasad Tetali. g -parking functions, acyclic orientations and spanning trees, 2010. [arXiv:0801.1114](#).
- 5 Tom Brylawski. The broken-circuit complex. *Trans. Amer. Math. Soc.*, 234(2):417–433, 1977.
- 6 Charles J Colbourn and William R Pulleyblank. Matroid steiner problems, the tutte polynomial and network reliability. *Journal of Combinatorial Theory, Series B*, 47(1):20–31, 1989. [doi:10.1016/0095-8956\(89\)90062-2](#).
- 7 Mary Cryan, Heng Guo, and Giorgos Mousa. Modified log-sobolev inequalities for strongly log-concave distributions, 2020. [arXiv:1903.06081](#).
- 8 Ewan Davies and Will Perkins. Approximately counting independent sets of a given size in bounded-degree graphs. *arXiv preprint*, 2021. [arXiv:2102.04984](#).
- 9 Yotam Dikstein, Irit Dinur, Yuval Filmus, and Prahladh Harsha. Boolean Function Analysis on High-Dimensional Expanders. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2018)*, volume 116 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 38:1–38:20, 2018.
- 10 I. Dinur and T. Kaufman. High dimensional expanders imply agreement expanders. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 974–985, 2017.
- 11 Andreas Galanis, Daniel Štefankovič, and Eric Vigoda. Inapproximability of the partition function for the antiferromagnetic ising and hard-core models. *Combinatorics, Probability and Computing*, 25(4):500–559, 2016.
- 12 I.M Gelfand, R.M Goresky, R.D MacPherson, and V.V Serganova. Combinatorial geometries, convex polyhedra, and schubert cells. *Advances in Mathematics*, 63(3):301–316, 1987. [doi:10.1016/0001-8708\(87\)90059-4](#).
- 13 Emeric Gioan and Michel Las Vergnas. The active bijection for graphs. *Advances in Applied Mathematics*, 104:165–236, 2019. [doi:10.1016/j.aam.2018.11.001](#).

- 14 Roy Gotlib and Tali Kaufman. No where to go but high: A perspective on high dimensional expanders, 2023. [arXiv:2304.10106](#).
- 15 Mark Jerrum and Alistair Sinclair. Approximating the permanent. *SIAM Journal on Computing*, 18(6):1149–1178, 1989. [doi:10.1137/0218077](#).
- 16 Richard M. Karp. *Reducibility among Combinatorial Problems*, pages 85–103. Springer US, Boston, MA, 1972. [doi:10.1007/978-1-4684-2001-2_9](#).
- 17 Tali Kaufman and Izhar Oppenheim. High order random walks: Beyond spectral gap. In *APPROX/RANDOM*, pages 47:1–47:17, 2018.
- 18 Milena Mihail and Umesh Vazirani. On the expansion of 0-1 polytopes. *preprint*, 3461, 1989.
- 19 Geoffrey Ramseier, Ashish Goel, and David Mazières. Liquidity in credit networks with constrained agents. In *Proceedings of The Web Conference 2020*. ACM, April 2020. [doi:10.1145/3366423.3380276](#).
- 20 Allan Sly. Computational transition at the uniqueness threshold. In *2010 IEEE 51st Annual Symposium on Foundations of Computer Science*, pages 287–296. IEEE, 2010.
- 21 Allan Sly and Nike Sun. Counting in two-spin models on d-regular graphs. *Annals of Probability*, 42(6):2383–2416, 2014.
- 22 Richard P. Stanley. Acyclic orientations of graphs. *Discrete Mathematics*, 5(2):171–178, 1973. [doi:10.1016/0012-365X\(73\)90108-8](#).
- 23 Richard P Stanley et al. An introduction to hyperplane arrangements. *Geometric combinatorics*, 13(389-496):24, 2004.

A Proof of Theorem 20

In this section we prove Theorem 20. We use a reduction from the problem of computing the partition function of the Hardcore model when the fugacity is above the critical threshold. Define $\#HC(\Delta, \lambda)$ as follows: given a Δ -regular graph $G = (V, E)$, compute the partition function $Z_G(\lambda) = \sum_I \lambda^{|I|}$, where the sum is taken over the family of independent sets $I \subseteq V$ of G . The critical threshold is defined as $\lambda_c(\Delta) := \frac{(\Delta-1)^{\Delta-1}}{(\Delta-2)^\Delta}$.

► **Theorem 23** ([20, 21, 11]). *The following holds for any fixed $\epsilon > 0$, integer $\Delta \geq 3$ and $\lambda > \lambda_c(\Delta)$: unless $NP=RP$, for any $\lambda > \lambda_c(\Delta)$ there is no polynomial-time algorithm for approximating $\#HC(\Delta, \lambda)$ up to a $1 + \epsilon$ multiplicative factor.*

We give a polynomial-time algorithm that given a $e^{\pm\epsilon/2}$ -approximation for $\#INDEP-SET-INC(7, \frac{2}{19})$ (see Definition 19), approximates $\#HC(7, \frac{2}{3})$ up to a $e^{\pm\epsilon}$ -multiplicative error. Since $\frac{2}{3} > \lambda_c(7) = \frac{6^6}{5^7} \geq 0.6$, this finishes the proof of Theorem 20. Our reduction is a modification of Theorem 16 in [8].

► **Theorem 24.** *There exists a polynomial-time algorithm that for any given $\epsilon \leq 1$, satisfies the following properties:*

1. *Given an instance $G = (V, E)$ of $\#HC(7, \frac{2}{3})$, the algorithm constructs an instance $G' = (V', E')$ of the problem $\#INDEP-SET-INC(7, \frac{2}{19})$ with size polynomial in $|G|$.*
2. *Given a $e^{\pm\epsilon/2}$ -multiplicative approximation to the number of independent sets of size $\lfloor \frac{2|V'|}{19} \rfloor$ of G' , a $e^{\pm\epsilon}$ -approximation of $Z_G(\frac{2}{3})$ can be computed in polynomial time.*

Proof. Given a 7-regular graph $G = (V, E)$, we define G' as the disjoint union of G with $r := \frac{c^2 n^2}{\epsilon}$ copies of the complete graph K_8 , where $n = |V|$, for some $c > 1$ that we choose later. For simplicity of notation, let $N := |V'| = n + 8r$, $\alpha := \frac{2}{19}$, $\lambda := \frac{2}{3}$. It is enough to show that G' is an instance of $\#INDEP-SET-INC(7, \frac{2}{19})$ and

$$e^{-\epsilon/2} \frac{i_{\lfloor \alpha N \rfloor}(G')}{\binom{r}{\lfloor \alpha N \rfloor} 8^{\lfloor \alpha N \rfloor}} \leq Z_G(\lambda) \leq e^{\epsilon/2} \frac{i_{\lfloor \alpha N \rfloor}(G')}{\binom{r}{\lfloor \alpha N \rfloor} 8^{\lfloor \alpha N \rfloor}}, \quad (9)$$

where as usual $i_k(G)$ is the number of independent sets of size k in G , and

$$\binom{r}{\lfloor \alpha N \rfloor} 8^{\lfloor \alpha N \rfloor} = i_{\lfloor \alpha N \rfloor}(rK_8).$$

Here, rK_8 is a shorthand for the graph which is a disjoint union of r copies of K_8 . We first show that Equation (9) holds. Note that

$$i_{\lfloor \alpha N \rfloor}(G') = \sum_{j=0}^n i_j(G) i_{\lfloor \alpha N \rfloor - j}(rK_8) = i_{\lfloor \alpha N \rfloor}(rK_8) \sum_{j=0}^n i_j(G) \frac{i_{\lfloor \alpha N \rfloor - j}(rK_8)}{i_{\lfloor \alpha N \rfloor}(rK_8)}.$$

Thus, to show Equation (9), it is enough to prove that for every $1 \leq j \leq n$,

$$e^{-\epsilon/2} \cdot \frac{i_{\lfloor \alpha N \rfloor - j}(rK_8)}{i_{\lfloor \alpha N \rfloor}(rK_8)} \leq \lambda^j \leq e^{\epsilon/2} \cdot \frac{i_{\lfloor \alpha N \rfloor - j}(rK_8)}{i_{\lfloor \alpha N \rfloor}(rK_8)}. \quad (10)$$

We can write

$$\frac{i_{\lfloor \alpha N \rfloor - j}(rK_8)}{i_{\lfloor \alpha N \rfloor}(rK_8)} = \frac{\binom{r}{\lfloor \alpha N \rfloor - j} 8^{\lfloor \alpha N \rfloor - j}}{\binom{r}{\lfloor \alpha N \rfloor} 8^{\lfloor \alpha N \rfloor}} = \frac{1}{8^j} \prod_{i=0}^{j-1} \frac{\lfloor \alpha N \rfloor - i}{r - \lfloor \alpha N \rfloor + j - i}. \quad (11)$$

To prove the upper bound, first note that

$$\frac{\alpha N}{r - \alpha N + j} \underset{\alpha N \geq 8\alpha r}{\geq} \frac{8\alpha r}{r(1 - 8\alpha) + n} \underset{\substack{n = \sqrt{\epsilon r}/c \\ \alpha = 2/19}}{=} \frac{16}{3} \left(\frac{1}{1 + \frac{19\sqrt{\epsilon}}{3c\sqrt{r}}} \right). \quad (12)$$

This implies that $\frac{\alpha N}{r - \alpha N + j} \geq 1$. So, $\frac{\alpha N}{r - \alpha N + j} \leq \frac{\lfloor \alpha N \rfloor - i}{r - \lfloor \alpha N \rfloor + j - i}$ for every $i < r - \lfloor \alpha N \rfloor + j$. Thus,

$$\begin{aligned} \frac{1}{8^j} \cdot \prod_{i=0}^{j-1} \frac{\lfloor \alpha N \rfloor - i}{r - \lfloor \alpha N \rfloor + j - i} &\geq \frac{1}{8^j} \cdot \left(\frac{\alpha N}{r - \alpha N + j} \right)^j \\ &\underset{\substack{\text{Equation (12)} \\ j \leq n = \sqrt{\epsilon r}/c}}{\geq} \frac{1}{8^j} \cdot \left(\frac{16}{3} \right)^j \left(\frac{1}{1 + \frac{19\sqrt{\epsilon}}{3c\sqrt{r}}} \right)^{\sqrt{\epsilon r}/c} \geq \left(\frac{2}{3} \right)^j e^{-\epsilon/2} = \lambda^j e^{-\epsilon/2}, \end{aligned}$$

for a large enough $c > 1$. Combining this with Equation (11), we get the upper bound in Equation (10).

To prove the lower bound, note that

$$\begin{aligned} \frac{1}{8^j} \cdot \prod_{i=0}^{j-1} \frac{\lfloor \alpha N \rfloor - i}{r - \lfloor \alpha N \rfloor + j - i} &\underset{j-i \geq 0}{\leq} \frac{1}{8^j} \cdot \left(\frac{\lfloor \alpha N \rfloor}{r - \lfloor \alpha N \rfloor} \right)^j \\ &\leq \frac{1}{8^j} \cdot \left(\frac{\frac{16r}{19} \left(1 + \frac{\sqrt{\epsilon}}{8c\sqrt{r}} \right)}{\frac{3r}{19} \left(1 - \frac{2\sqrt{\epsilon}}{3c\sqrt{r}} \right)} \right)^j \\ &\leq \left(\frac{2}{3} \right)^j e^{\epsilon/2} = \lambda^j \cdot e^{\epsilon/2}, \end{aligned}$$

for a large enough $c > 1$. Combining this with Equation (11), the lower bound in Equation (10), thus (9) follows.

It remains to show that G' is an instance of $\#$ INDEP-SET-INC $(7, \frac{2}{19})$, i.e. $i_k(G') \leq i_{\lfloor \alpha N \rfloor}(G')$ for any $k < \lfloor \alpha N \rfloor$.

40:14 On Optimization and Counting of Non-Broken Bases of Matroids

For any $k < \lfloor \alpha N \rfloor$, and any independent set S in the original graph G , let $T_{S,k}$ be the set of all independent sets of size k of G' whose intersection with the vertices of G is S . It is enough to show that there exists a constant n_0 such that if $n \geq n_0$, then we have $|T_{S,k}| \leq |T'_{S, \lfloor \alpha N \rfloor}|$ for every independent set $S \subseteq V$ of G and $k < \lfloor \alpha N \rfloor$. We prove a stronger statement that there exists a constant n_0 such that if $n \geq n_0$, then for any fixed independent set $S \subseteq V$, $|T_{S,k}|$ is increasing as a function of k for all $k \leq \lfloor \alpha N \rfloor$. It is enough to show that $\frac{|T_{S,k}|}{|T_{S,k-1}|} \geq 1$ for any $|S| \leq k \leq \alpha N$. Note that $|T_{S,k}| = \binom{r}{k-|S|} 8^{k-|S|}$. So we have

$$\frac{|T_{S,k}|}{|T_{S,k-1}|} = \frac{\binom{r}{k-|S|} 8^{k-|S|}}{\binom{r}{k-1-|S|} 8^{k-1-|S|}} = 8 \cdot \frac{r-k+|S|+1}{k-|S|} \geq 8 \cdot \frac{r-k}{k} \geq 8 \frac{\frac{3r}{19} - n}{\frac{16r}{19} + n},$$

where the last inequality comes from the fact that $k \leq \frac{2N}{19} = \frac{2}{19}(8r+n) \leq \frac{16r}{19} + n$. But since $r = \frac{c^2 n^2}{\epsilon}$, there is a constant n_0 such that for $n \geq n_0$, we have $\frac{\frac{3r}{19} - n}{\frac{16r}{19} + n} \geq \frac{1}{8}$. This shows that $\frac{|T_{S,k}|}{|T_{S,k-1}|} \geq 1$, which finishes the proof. \blacktriangleleft